

155. *Tettigonia fuscolineella*, sp. n. (Tab. XIX. fig. 25.)

Brevis, robusta; capite lato, perbrevis, rotundato, oculis prominulis, rufo-brunneo, fusco-variegato, lineâ ante ocellos et basi testaceis; pronoto convexo, nitido, antice rufo-brunneo, postice olivaceo, basi anguste testaceâ lineâ obscuriori ante colorem testaceum instructo; scutello rufo-brunneo; tegminibus olivaceis, fusco regulariter lineatis, marginibus anguste dilutioribus, apice subhyalino; abdomine rufo-brunneo; sterno pedibusque brunneis vel rufo-brunneis.

A short, small, robust species; head very short, broad, and rounded, the eyes prominent, reddish-brown, more or less infusate, with the base narrowly testaceous and a narrow testaceous line before the ocelli; pronotum convex and shining, olivaceous, with the anterior margin more or less broadly reddish-brown, the base narrowly testaceous; before the testaceous space is a dark brown line, and there is usually a testaceous line behind the reddish-brown portion in front; scutellum reddish-brown; tegmina olivaceous, with very regular fuscous lines, the apex subhyaline; wings smoky; abdomen reddish-brown; sternum and legs brown or red-brown: in some specimens the colour is much brighter than in others.

Long. 5-6 millim.; lat. $1\frac{3}{4}$ millim.

Hab. MEXICO, Atoyac, Vera Cruz (*H. H. Smith*), Orizaba (*Sallé*, *F. D. Godman*, & *H. H. Smith*); PANAMA, Bugaba (*Champion*).

An example from Atoyac is figured.

Var. fronteræ, n.

Cum formâ typicâ congruens, sed tegminibus fuscatis haud vel vix lineatis, suturâ clavali dilutiori facile distinguenda.

Closely allied to the type-form, but slightly smaller, with the tegmina of a uniform dark velvety-brown or blackish-fuscous colour, without distinct lines, the claval suture being somewhat lighter.

Long. 5 millim.; lat. $1\frac{1}{2}$ millim.

Hab. MEXICO, Frontera in Tabasco (*H. H. Smith*).

156. *Tettigonia psittacella*, sp. n. (Tab. XIX. fig. 26.)

T. fuscolineellæ affinis, sed minor, viridis vel ochraceo-viridis; capite lato, perbrevis, oculis prominulis, nigro, basi et lineâ ante ocellos flavis; pronoto viridi, antice et postice anguste flavescente, lineâ nigrâ ante colorem flavum ad basim instructo; scutello flavescente; tegminibus virescentibus, concoloribus, apicibus subhyalinis; abdomine nigro; corpore subtus plerumque nigro; pedibus flavis vel flavo-testaceis.

Smaller than *T. fuscolineella*, to which it is closely allied; green or olivaceous-green, the head black, with a yellow line before and behind the ocelli; the pronotum narrowly yellow in front and behind, and with a black line before the yellow base; the scutellum yellow, sometimes with a black patch at the base, and the tegmina unicolorous, without lines, the apex being subhyaline with a small fuscous spot at the tip; head very short, broad, rounded, eyes prominent; pronotum slightly broader at the base than the head with the eyes; scutellum rather small; tegmina narrowed to the apex; abdomen fuscous; underside for the most part fuscous; legs yellow or yellow-testaceous.

Long. 4-5 millim.; lat. $1-1\frac{1}{3}$ millim.

Hab. MEXICO, Orizaba (*Sallé*), Atoyac in Vera Cruz, Teapa in Tabasco (*H. H. Smith*); GUATEMALA, Pantaleon (*Champion*); PANAMA, Bugaba (*Champion*).

This and the preceding species will probably form the types of a new genus, as they differ considerably in the shape of the head from the ordinary forms of the genus.

We figure a specimen from Teapa.