

Genera.	Net Total of Species [] = unde- scribed in Central America.	Distribution in								
		Central America								and
		Mexico.	British Honduras.	Guatemala.	Salvador.	Honduras.	Nicaragua.	Costa Rica.	Panama.	Other parts of the World.
Lachlania	1	1	1	Cuba.
Homœoneuria ..	1	1	
Euthyplocia	1	1	..	1	1	Tropical South America.
Campsurus	2+[1]	1	[1]	1	Texas, Brazil.
Hexagenia	1	1	Canada to Brazil, India.
Choroterpes	2	1	..	1	Arizona, Europe, Te- nasserim.
Thraulius	6+[3]	3+[1]	..	1+[2]	1	4	Colombia, Europe, Indo - Malayan Region.
Gen. —	[1]	[1]						
Ephemerella	[1]	[1]	Northern Temperate Region.
Tricorythus	1	1	Egypt, S. Africa, Ma- layan Region.
Leptohyphes	1	1	Argentine Republic.
Baëtis.....	1+[3]	[3]	..	1	1	..	Nearly everywhere.
Centroptilum	[1]	[1]	Cuba, N. America, Europe.
Callibætis	2+[1]	1+[1]	..	2	N. and S. America, Australia.
Chirotonetes	[1]	[1]	..	[1]	N. America, Europe, Indo-MalayanRegion, Japan.
Gen. —	[1]	[1]								

The unnamed genus following *Thraulius* in the above table of geographical distribution works out with *Adenophlebia*; their differences are set forth below in the description. The other unnamed one, next to *Chirotonetes*, is related to *Cinygma*, a N.-American genus.

LACHLANIA.

Lachlania, Hagen, Proc. Bost. Soc. Nat. Hist. 1868, pp. 372-374 (woodcut).

This genus contains a Cuban species in addition to the one from Central America.