

Guadalupe [*Hay*: 1 ♂, 1 ♀] (*U. S. N. M.*), Mexico City (*H. H. Smith, Schumann, Deam*: 12 ♂, 17 ♀) [3 ♂, 6 ♀], Chapultepec [4 ♂], Tacubaya [1 ♂] (*Barrett, colls. Adams, P. P. C.*), Tlalpam (*Tower, coll. P. P. C.*: 1 ♂, 1 ♀).—BERMUDAS^{5 6 13 16}, Hamilton I. (*Snyder, A. N. S.*: 1 ♀); BAHAMAS⁸.

The Mexican examples were taken from June to September, those from San José del Cabo and Hamilton I. in October.

Prof. Blatchley records¹⁷ the Asilid, *Proctacanthus philadelphicus*, attacking this species at Ormonde, Florida.

SUPPLEMENT.

HETÆRINA (p. 19).

Heterina, Needham, Bull. 68, N. Y. St. Mus. p. 220 (1903).

The supplementary material requires the following changes and additions to the Key to Species, *anteà*, pp. 20–23:—

§ I. to be united with § II.

Under “*Males*” of § II.: rubric *e*, after “pale brown” insert “(old males of *cruentata* may have the labrum almost or entirely black)”; add a new rubric:—

“*eee*. Labrum black, usually with a yellow spot each side; tips of all the wings with a brown spot 1. *fuscoguttata*.”

Under “*Females*” of § II.: rubric *f*, after “obscure in colour” insert “(some females of *capitalis* will fall here, but see page 347, *postea*)”; under *dd*, insert:—

“*dd* 1. Metasternum with no circular black border. (Includes *g* and *gg* as before, 7. *titia* and 4. *americana*; *americana* usually has the tibiae pale superiorly, *titia* not so.)

dd 2. Metasternum with a circular blackish border, slightly interrupted anteriorly.

k. Size larger (length, abd. 33–35, hind wing 27·5–29·5 mm.), 21–26 antecubitals on the front wings, wings with no stigma, venation at base usually reddish. 1. *fuscoguttata*.