

Heterina americana.

	Chihuahua City.				Baños de Santa Rosalia.			Escuinapa.			Gualan.		Zacapa.		Amatitlan.		Fscuintla.
	7 ♂.	1 ♂.	1 ♂.	1 ♀.	1 ♂.	2 ♂.	2 ♀.	1 ♂.	2 ♀.	2 ♀.	♂ ♂.	♀ ♀.	♂ ♂.	♀ ♀.	♂ ♂.	♀ ♀.	
Number of cells surmounted by stigma of front wings	1+ to 2+	less than 1	*	*	1+	*	1+ to 2+	1 to less	*	*	1 to 2+	1+ to 2+	1+ to 2+	1+ to 2+	1+ to 2+	1+ to 2+	2 ♂.
Number of cells surmounted by stigma of hind wings	1+ to 2+	*	*	*	1+ to 2+	*	1+	less than 1 to 0	*	*	1 to 2+	1+ to 2+	1+ to 2+	1+ to 2+	1+	1+	1+ to 2+
Fractional part of distance from base to nodus to which red of front wings reaches, ♂5-.67	.64	.73	..	.5	.3-.72	..	.7	.73	..	.45-.55	..	.45-.5	..	.4-.45	..	.5
Figs. of Tab. II. which the ♂ appendages most resemble	6, 14 15	12	15	..	15	12	..	12	12	..	6, 14	..	6, 14	..	13	..	6, 15
Length of abdomen in mm.	34-36.5	34.5	34	29	35	30-31	29.5-31	?	36	28.5	32-34.5	26.5-28	31-34	25.5-28	30-32	26	32-33.5
Length of hind wing in mm.	25-26	25	26	26	25	21.5-22	26-26.5	24	26	26-26.5	23-25	24-25	23-24	23-25	22.5-23.5	23-24	23-23.5

* Pterostigma absent.

† Stigma represented by a clouding around a cross-vein.