

12. *Anaplecta albomarginata*, sp. n.

♀. *A. decipienti* simillima, eadem pictura, abdomine tamen basi testaceo. Differt ab illa:—Formis gracilioribus; elytris sensim angustioribus, apice regulariter rotundatis, margine costali minus areuato; campo discoidali tantum apice reticulato. Alæ hyalinæ, venis costalibus fulvescentibus, campo appendiculari sat grandi, longiore quam latiore, hyalino, margine basali subrecto. Ultimum segmentum ventrale apice haud compressum, sed rotundato-tumidum, fissum. (Venæ campi discoidalis in elytro sinistro longitudinales; vena media in elytro dextro oblique tiramosa.)

Long. 5; elytr. 5, pronot. 1.5, latit. pronot. 2 millim.

*Hab.* MEXICO, Tabasco (*H. H. Smith*).

13. *Anaplecta parvipennis*, sp. n.

Minuta, statura *A. grandipennis*, nigro-castanea, antennis, pedibus et cercis testaceis. Pronoti margines laterales hyalino-pellucentes. Elytra abdomen haud superantia, margine costali albo, apicali testaceo, apice rotundato, margine costali arcuato, margine suturali tantum apice arcuato. Alæ minutæ, fusco-nebulosæ, ad marginem costalem fusciores; campo appendiculari brevi, rotundato, brevior quam latiore, margine basali vix angulato. Area medio-discoidalis venula transversa unica angulata, venam longitudinalem sat brevem, obliquam, emittens. Lamina supra-analis transversa.—*Variat.* Alis vix infuscatis.

Long. 4.8; elytr. 3.4, pronot. 1.3, latit. pronot. 2.2 millim.

*Hab.* MEXICO, Tabasco (*H. H. Smith*).

Of the same facies as *A. azteca*, *A. replicata*, *A. grandipennis*, *A. decipiens*, *A. albomarginata*, and *A. fallax*, but distinct from all these (*A. azteca* excepted) on account of the wings being one half shorter and the elytra narrower. Smaller than *A. decipiens*, with the elytra more rounded at the tip and not reticulated, and the wings with shorter appendicular field. A little larger than *A. replicata* and *A. grandipennis*, with very differently formed wings (comp. synopsis specierum II. 2, *anteà*, p. 21). Distinguished from *A. albomarginata* by the shorter elytra, infuscated wings with shorter appendicular field, &c. It is quite similar to *A. fallax*, only the elytra are a little shorter, not longer than the abdomen, and the wings one half smaller, a little more infuscated, with the appendicular field a little shorter and the longitudinal venula of the end of the medio-discoidal area shorter and oblique. *A. parvipennis* is remarkable for its small hind wings—when extended reaching scarcely to the end of the abdomen—being scarcely longer than the elytra. It is the only known species possessing this character.

14. *Anaplecta domestica*, sp. n.

♀. Minuta, statura *A. fallacis*; crassiuscula, rufo-castanea vel rufescens, subtus rufo-testacea; pedibus plus minus testaceis. Antennæ rufo-testaceæ. Pronotum leviter bi-impressum, marginibus lateralibus totis hyalino-testaceis. Elytra abdomen vix superantia, apice subacuminato-rotundato, margine costali arcuato, suturali recto; margine costali saltem basi testaceo; campi discoidalis venis longitudinalibus. Alæ hyalinæ, venis testaceis; area medio-discoidali venulis transversis 1-2 vel ante apicem venula unica obliqua, ut in *A. grandipenne*. Campus appendicularis hyalinus, rotundatus, æque longus ac latus, mediocris, plus quam 3<sup>a</sup> parte longitudinis reliqui alæ æqualis. Lamina supra-analis brevis, transversa.

Long. 4.6; elytr. 3.8, pronot. 1.75, latit. pronot. 2.25 millim.

*Hab.* GUATEMALA, in the city (*Champion*).

A species very like *A. alaris*, but with the elytra more rounded at the tip, with the