

b. Testacea; pronoti discus fuscus, circumcirca testaceo-marginatus, per vittam flavidam divisus.—16. *divisa*, sp. n.

bb. Rufo-testacea; pronotum corneum, disco fasciis 2 arcuatis fuscis.—[17. *bolliana*, sp. n.]

11. Femora anteriora margine infero-interno tota longitudine spinoso, spinis partis distalis minoribus, confertis, distinctissimis. (Species mundi antiqui.)—(ISCHNOPTERA.)—18. *blattoides*, Sss. (Cf. *I. ectobioides*, Sss.; *I. bocagei*, Bol.; *I. vitticollis*, Br., &c.)

Conspectus specierum secundum alam ordinatarum.

a. Alarum area medio-discoidalis areae medio-ulnari æquilata.—1. *inæqualis*; [2. *uhleriana*]; 6. *ignobilis*; 7. *nahua*; (8. *inca*); 13. *parvula*; 16. *divisa*; 18. *blattoides*.

aa. Alarum area medio-discoidalis quam area medio-ulnaris evidenter latior.—3. *mexicana*; 4. *consobrina*; 5. *conformis*; 8. *inca*; 9. *castanea*; 10. *azteca*; 11. *tolteca*; 12. *nana*; 14. *annulicornis*; 15. *couloniana*; [17. *bolliana*].

1. Ischnoptera inæqualis, sp. n. (Tab. VI. figg. 14–17.)

- ♂. Fusca vel fusco-ferruginea, antennis brunneis. Caput facie testacea, in medio et vertice nigris. Pronotum ellipticum, impressionibus distinctis, disco convexiusculo, lævigato, toto nigro vel fusco vel castaneo, marginibus lateralibus testaceis, translucidis, fere sulfureis. Elytra fusca vel rufescentia, area basali marginis antici hyalina. Alæ subvitreæ, margine anteriore et apicali latiuscule infuscato. Vena ulnaris 6–7-ramosa; præterea ramo incompleto, alterisque rudimentariis, venam dividenter haud attingentibus.
- ♀. Pronotum corneum, parabolicum, margine anteriore et laterali semi-elliptico, posteriore transverso, lato, subangulato, angulis lateralibus leviter rotundatis. Discus convexiusculus, impressionibus nullis, fusco-niger, marginibus lateralibus flavidis. Elytra abbreviata, primum abdominis segmentum tegentia, fusco-nigra, venis distinctis, apice late rotundata, area costali flavida. Alæ rudimentariæ, flavicantes, segmentum mediale haud superantes, apice fuscae vel maculosæ.
- ♀. Long. 16; pronot. 5, latit. pronot. 6·5, long. elytr. 8, latit. elytr. 5·25 millim.
- ♂. Long. 21; pronot. 5·2, latit. pronot. 6·3, long. elytr. 23, latit. elytr. 6·8 millim.

Hab. NORTH AMERICA, Texas.—NORTH MEXICO (*Mus. Genavense*).

This species differs from *I. coulöniana* (♂) in its pronotum being more coriaceous and quite black, except its lateral margins.

[2. Ischnoptera uhleriana, Sauss. (Tab. III. figg. 21–23.)

Ischnoptera uhleriana, Sauss. Revue et Mag. Zool. xiv. p. 169 (1862); Mém. Mex., Blatt. p. 82. 23 (♂); Miss. Scient. Mex., Orthopt. p. 55. 2 (♂).

- ♂. Pallide testacea, fulva. Pronotum ellipticum, utrinque subdeflexum, circumcirca pallidius; impressionibus disci parum profundis, obsoletis. Elytra fulvescentia. Alæ subvitreæ, venis testaceis, venulis costalibus medii campi marginalis incrassatis, brunneis; vena ulnari ramis apicalibus 5–6, alterisque abbreviatis 2–3. Lamina supra-analis trigonalis. (Var. *fulvescens*.)
- ♀. Fuscescens, capite, pronoto, elytris pedibusque testaceis. Caput flavidum, facie media brunescens. Pronotum quam in ♂ minus ellipticum, subcorneum, impressionibus obliquis disci obsoletissimis; margine posteriore minus arcuato, disco rufo-brunneo. Elytra abbreviata ad 3^m abdominis segmentum extensa, fulvo-testacea, apice late rotundata, margine exteriori arcuato, apice fere regulariter rotundato, campo costali translucido, venis brunneis. Alæ breviores, primum abdominis segmentum tegentes. Abdomen superne et inferne fuscum; segmentis margine laterali et posteriore testaceo. Lamina supra-analis trigonalis. Cerci testacei, basi et subtus fusi.
- ♀. Long. 11·7; elytr. 6, latit. elytr. 3·6, pronot. 3·5, latit. pronot. 4·75 millim.
- ♂. Long. 17; elytr. 17·8, latit. elytr. 5·2, pronot. 4, latit. pronot. 5·3 millim.

Hab. NORTH AMERICA, Georgia, Texas, New Mexico.]