

*Hab.* MEXICO, Teapa in Tabasco (*H. H. Smith*).

Very close to *B. azteca*, but somewhat smaller; the lamina supra-analis very slightly split at the end; the wings rather shorter; the vena mediastina forming only two branches.

### 7. *Blatta totonaca*, Sauss.

*Blatta totonaca*, Sauss. Rev. et Mag. de Zool. 1869, p. 165; Mém. Mex., Blatt. p. 101. 40; Miss. Scient. Mex., Orthopt. p. 37. 13, t. 1. fig. 23.

*Hab.* MEXICO, Orizaba (*Mus. Genavense*).

### [8. *Blatta supellectilium*, Serv.

*Phyllodromia supellectilium*, Serv. Orth. p. 114. 44; Brunn. de W. Syst. Blatt. p. 98. 11 (1865); Savigny, Descript. de l'Égypte, Orth. t. 2. fig. 19; Sauss. Miss. Scient. Mex., Orthopt. p. 39. 14.

*Blatta cubensis*, Sauss. Rev. et Mag. de Zool. 1862, p. 166; Mém. Mex., Blatt. p. 108. 46, figg. 14, 15 (♀ ♂).

*Phyllodromia cubensis*, Brunn. de W. Syst. Blatt. p. 109. 29.

*Hab.* BRAZIL; ANTILLES, Cuba (*Saussure*).—TROPICAL AFRICA (*Mus. Genavense*); MAURITIUS.]

### 9. *Blatta fraterna*, sp. n.

♂. Sat minuta, testacea. Oculi remoti. Frons inter illos vitta, necnon linea transversa supra-antennali, fuscis. Facies punctis nonnullis fuscis. Pronotum ellipticum, margine posteriore vix arcuato, disco flavicante, punctis rarioribus fuscis. Elytra sat brevia, abdomen superantia. Alæ vitreæ, venis fusco-testaceis; margine anteriore in medio fusco-umbrato; venis costalibus paucis, vix incrassatis. Pedes testacei. Lamina supra-analis trapezina laminam infra-genitalem superans, obtusangulatim incisa, leviter late bilobata. Cerci longissimi, apice articulis valde discretis, ultimo apice fusco. Lamina infra-genitalis margine posteriore lato, leviter arcuato, subinciso, stylis invicem valde remotis ad angulos laterales positus. Pedes testacei.

Long. 11.5; elytr. 11, latit. elytr. 4, pronot. 4, latit. pronot. 5.25 millim.

*Hab.* NICARAGUA, Chontales (*Janson*).

A species much resembling *B. nahua* in the form of its anal parts, but smaller; the eyes more distant, not strongly convergent from above; the elytra much shorter; and the wings with much less numerous costal veins.

### 10. *Blatta dilatata*, Sauss.

*Blatta dilatata*, Sauss. Rev. et Mag. de Zool. 1868, p. 98. 6 (♀); Miss. Scient. Mex., Orthopt. p. 32. 7, t. 1. fig. 19 (♀) (nec ♂, ad *Bl. vitream* pertinens).

Lamina supra-analis ♀ trigonalis, sulco divisa, ultimum segmentum ventrale haud superans. Long. 10; elytr. 9.75, pronot. 3.4, latit. pronot. 5.2 millim.

*Hab.* MEXICO, Orizaba.