

11. **Blatta vitrea**, Brunn.

*Phyllodromia vitrea*, Brunn. de W. Syst. Blatt. p. 109. 28 (♂).

*Blatta vitrea*, Sauss. Miss. Scient. Mex., Orthopt. p. 30. 6, t. 1. fig. 18 (♂).

*Blatta dilatata*, Sauss. Rev. et Mag. de Zool. 1868, p. 98. 6 (♂) (nec ♀, cfr. no. 10).

*Hab.* MEXICO, Eastern Cordillera, Sierra de Moyoapam (*Saussure*).

12. **Blatta orizabæ**, Sauss.

*Blatta orizabæ*, Sauss. Rev. et Mag. de Zool. 1868, p. 355. 8 (♂ ♀); Miss. Scient. Mex., Orthopt. p. 39. 15, t. 1. figg. 24 *a, b*.

*Hab.* MEXICO, Eastern Cordillera, Orizaba (*Saussure*).

13. **Blatta acolhua**, Sauss.

*Blatta acolhua*, Sauss. Rev. et Mag. de Zool. 1868, p. 99. 8 (♀); Miss. Scient. Mex., Orthopt. p. 40. 16, t. 1. fig. 25 *a*.

*Hab.* MEXICO, Eastern Cordillera (*Saussure*), Vera Cruz (*Schumann*), Teapa in Tabasco (*H. H. Smith*).

14. **Blatta maya**, sp. n.

Pallide testacea, leviter rufescens; pronoti disco rufo-flavido vel obsolete brunneo-irrorato. Elytra mediocriter longa, leviter fulvescentia. Alæ amplæ, venulis transversis campi antici fuscis; venis costalibus apice leviter incrassatis, haud coloratis. Lamina supra-analis ♀ trigonalis, basi carinata, apice breviter fissa, obtuse bidentula, ♂ transversa, margine postico arcuato. Lamina infra-genitalis ♂ in medio producta et impressa, truncata, angulis rotundatis, stylis brevibus, apicalibus, invicem propinquis instructa.

Long. 12; elytr. 12.5, pronot. 4, latit. pronot. 5 millim.

*Hab.* MEXICO, Temax in North Yucatan (*Gaumer*).

15. **Blatta zapoteca**, Sauss.

*Blatta zapoteca*, Sauss. Rev. et Mag. de Zool. 1862, p. 166; Mém. Mex., Blatt. p. 105. 44; 3 s. Scient. Mex., Orthopt. p. 34. 10, t. 1. fig. 21.

♀. Media, fulvo-testacea; abdomine fusco vel infuscato. Antennæ basi fuscae. Caput nigrum, vertice inter oculos flavido, lineolis 4 fuscis; ocellis, macula inter antennis alteraque clypei, flavis. Oculi a supero visi convergentes, invicem remoti. Pronotum testaceum, ellipticum, margine postico vix arcuato, disco fusco-marmorato, amœne multipicturato, lateribus frequenter fusco-punctatis, margine laterali subtiliter fusco. Elytra modice longa, abdomen superantia, fulvescentia, vena humerali et discoidali basi fuscis. Alæ subvitreæ, venis fusco-testaceis; margine anteriore et apice fulvescentibus; venis costalibus apice elongato-incrassatis. Pedes testacei, ad spinas fusco-punctati. Femora superne et interdum subtus fusco-marginata. Tibiæ et articuli tarsorum apice fuscis. Lamina supra-analis trigonalis, carinata, apice angulato-incisa. Segmentum ventrale haud superans. Cerci modice longi.

*Var.* Caput testaceum, fusco-maculatum. Abdomen piceum vel testaceum. Femora vix fusco-marginata.

Long. 14.5; elytr. 13, pronot. 4, latit. pronot. 6.25 millim.

*Hab.* MEXICO (*Saussure*); GUATEMALA, Capetillo (*Champion*).

A very distinct species, remarkable on account of the curious brown picturing of its pronotum, marmorated with regular drawings, sometimes imitating flowers, or, if more brown is present, with yellow spots.