

11. Elytra abbreviata, squamiformia vel nulla. Pronotum parabolicum, postice transversum.
2. Caput prominulum. Femora anteriora subtus margine postico inermia. ♂♂ complete alati. Abdominis margines haud serrati. Lamina supra-analis ultra segmentum ventrale vix prominulum, transversa, arcuata, vel vix incisa.—*CALOLAMPRA*, Sss.
22. Caput obtectum. Corpus depressum, ellipticum.
3. Lamina supra-analis transversa, arcuata, subincisa, vix prominula. Corpus nitidum.
4. Femora anteriora subtus margine antico piloso, postico spina unica. Elytra ♀♂ truncata. Corpus gracile; abdominis margines haud serrati. (*Ep. liturata* ♂, De Haan*).—[*COMPSOLAMPRA*, Sss.†]
44. Femora anteriora in utroque margine spinosa. Elytra squamiformia. Corpus late ellipticum, abdominis margines serrati.—[*OPISTHOPLATIA*, Br.]
33. Lamina supra-analis prominula, trigonalis, incisa. Femora anteriora in utroque margine spinosa. Corpus rugulosum; pronotum cucullatum. Abdomen valde serratum. Elytra squamiformia vel nulla.—*RHICNODA*, Br.

PARATROPA, Serv.

Paratropes, Serville, Orthopt. p. 117; Saussure, Miss. Scient. Mex., Orthopt. p. 73.

Phoraspis, Burmeister, Handbuch, ii. p. 492.

Paratropa, Brunner de W. Syst. Blatt. p. 148.

Synopsis specierum.

- a.* Alæ aurantiæ, fusco-marginatæ.
- b.* Pronotum anterius vitta flava, parallela vel in medio vix coarctata.—[*subsericea*, Sauss. (Guiana)]; [*elegans*, B. (Guiana)].
- bb.* Pronotum anterius vitta flava in medio valde coarctata vel interrupta.
- c.* Vitta haud interrupta.
- d.* Pronoti margo posterior immaculatus.—1. *mexicana*, Br.; [*æquatorialis*, Sauss. (Peru)].
- dd.* Pronoti margo posterior flavo-bimaculatus.—[2. *phalerata*, Er.]
- cc.* Pronotum anterius flavo-bimaculatum.—3. *bilunata*, sp. n.
- aa.* Alæ totæ fuscae. Pronotum ut in *P. mexicana* pictum.—4. *biolleyi*, sp. n.

1. *Paratropa mexicana*, Brunn.

Paratropes lycus, var. ♀, Sauss. Mém. Mex., Blatt. p. 60.

Paratropa mexicana, Brunn. de W. Syst. Blatt. p. 151. 2, fig. 15 (♀); Sauss. Miss. Scient. Mex., Orthopt. p. 73. 2.

Hab. MEXICO, Eastern Cordillera, Orizaba (*Mus. Genavense, ex coll. Guérin*).

[2. *Paratropa phalerata*, Erichs.

Blatta (*Nyctibora*) *phalerata*, Erichson, in Schomb. Fauna und Flora v. Britisch-Guiana, p. 580 (1848).

* Nec ♀.—*Epil. quadrata*, Sauss. Mém. Orth. 4^e fasc. p. 129, ♀♂.

† Sauss. Societas Entomologica, 1893, p. 58. Asiatic type mentioned here to fix the differences of *Calolampira*.