

♂. Alarum campus anterior campum posteriorem leviter superans. Lamina supra-analis minus producta, parum incisa, a cercis superata. Vertex inter oculos eorum latitudine æqualis.

♀. Long. 31; elytr. 26·5, pronot. 6·75, latit. pronot. 8·5 millim.

Hab. GUATEMALA, Vera Paz (*Champion*).—GUIANA, Cayenne.

This species has the pronotum very finely punctured with brown, as in *E. caraibea*. *E. caraibea*, however, is much smaller than *E. conspersa*, and the head is not densely punctured with brown in the same way.

Var. ? ♂. Antennæ, caput, pedes et cerci nigra (anne fortuito?). Elytra tenuissime rufo-tessellata, fere ut in *E. caraibea*, sed obsoletius. Long. 23; elytr. 25, pronot. 5·75, latit. pronot. 8 millim.

Hab. PANAMA, near the city (*Champion*).

It is somewhat doubtful whether this variety (♂) belongs to *E. conspersa*, the eyes being less distant, and the elytra having a different sort of fine maculation. Its black head and legs may be the result of discoloration in the dried specimens.

6. *Epilampra azteca*, Sauss.

Epilampra azteca, Sauss. Rev. et Mag. de Zool. 1868, p. 356. 12 (♂); Miss. Scient. Mex., Orthopt. p. 82. 5, t. 2. fig. 45 (pars).

♀. Flavo-testacea, facie punctis 2 vel 4 et linea media fuscis. Frons cum pictura frequenter lineas 2 ad occipitem emittente necnon maculis 2 ad oculos, fuscis. Oculi quam ocelli æque remoti. Pronotum paulo latius ac brevius quam in *E. caraibea* et *E. latifrons*; postice bisinuat, utrinque grosse sparse fusco-punctatum; disco pictura magna marmorata fusca, distinctissima, haud fusco-punctulata, notato. Elytra fulvescentia, omnino crasse, remote, fusco-punctata; linea in vena humerali elongata fusca ad basin evanida, maculaque basali campi analis ad venam analem fusca. Alæ leviter infuscatæ, margine costali fulvescente. Campus anterior angustus, apice leviter truncatus vel rotundatus, campum posticum vix superans. Incisura apicalis inter campos nulla; apex alæ inter campum anticum et posticum aream membranaceam intercalatam angustam præbens. Abdomen supra testaceo-flavum, segmentis posterioribus utrinque castaneo-umbratis, angulis leviter prominulis, subacutis, 7° segmento leviter prominulo, angulis leviter productis, rotundatis. Venter castaneus, limbo et basi luteis, sparse fusco-punctatis. Lamina supra-analis ♀ longiuscula, paulum prominula, retro attenuata, rotundata, leviter incisa, bilobata, basi fusca. Ultimum segmentum ventrale inter cercos productum, margine valde arcuato, subtruncato. Cerci modice longi, testacei, laminam supra-analem haud longe superantes, apice anguste attenuato, haud nigro.

Long. 18; elytr. 17·5, pronot. 4·25, latit. pronot. 6·5 millim.

Hab. MEXICO, Eastern Cordillera (*Saussure*), Teapa in Tabasco (*H. H. Smith*).

Saussure formerly included specimens from Cuba and others from South America under *E. azteca*; these are here separated under the names *E. caraibea* and *E. latifrons* respectively.

[7. *Epilampra caraibea*, sp. n.

E. aztecæ simillima, at minus fusco-picturata. Caput ubique sparse fusco-punctatum, fronte infra antennas vitta angusta transversa flavida; vertice inter oculos nigro, saltem pictura nigra. Caput superne flavidum, occipite fusco-binotato. Pronotum omnino subtilissime fusco-punctulatum, in disco densissime, utrinque sparsius, necnon ad margines laterales punctis majoribus fuscis; pictura lyrata fusca nulla.

BIOL. CENTR.-AMER., Orthopt., November 1893.

kk