

In this species the marking of the pronotum is intermediate between that of *E. azteca* and that of *E. caraibea*; the markings of the elytra are also intermediate, being more finely and less irregularly punctured than in *E. azteca*, but more irregularly than in *E. caraibea*; the colour of the abdomen on the upperside is like that of *E. caraibea*.]

CALOLAMPRA, Sauss.

Calolampra, Saussure, Societas Entomologica, 1893, p. 58.

Generi *Epilampræ* simillimum; differt tamen feminis apteris vel elytris valde abbreviatis squamiformibus; marium elytris complete explicatis vel atrophicis. Femora anteriora subtus margine postico inermi, tantum spina apicali instructo, margine antico inermi vel basi spinoso, apice spinis 1 vel 2 majoribus.

Synopsis specierum.

a. Femora anteriora spinosa. Elytra ♀ rudimentalia vel nulla.

b. Femora subtus margine postico inermi; femora anteriora margine antico basi spinis normalibus, apice spinulis minimis piliformibus instructo. (Blattinæ, Type 2, *Platamodes*, anteà p. 30). Elytra ♀ rudimentalia, ♂ completa. Tarsi graciles, metatarso postico quam reliqui articuli computati haud brevior.—[*gracilis*, Br. (Australia)]; [*heusseriana* *, Sss.].

bb. Femora 2^a, 3^a subtus in utroque margine spinosa. Tarsi breves, crassiusculi. Feminae apteræ. Mares?—1. *brevitarsis*, Sss.

aa. Femora anteriora in utroque margine mutica; posteriora saltem in margine postico rare spinosa. Elytra marium completa.—2. *bispinosa*, Sss.

1. *Calolampra brevitarsis*, Sauss. (Tab. IV. fig. 39.)

Calolampra brevitarsis, Sauss. Societas Entomologica, 1893, p. 58.

♀. Gracilis, depressa, fusco-nigra, nitida. Caput prominulum, vertice rotundato, ore et ocellis testaceis. Antennæ crassiusculæ, fusco-testaceæ. Pronotum corneum, parabolicum, antice truncatum, margine posteriore parum arcuato. Mesonotum et metanotum angulis posticis nullomodo productis; marginibus lateralibus mesonoti arcuatis. Thorax coriaceo-punctulatus. Pedes fusco-testacei, sat robusti. Femora anteriora subtus margine antico piloso-spinuloso, ad basin spina 1, apice 2. Femora intermedia et posteriora remote spinosa, spinis marginis antici brevibus. Tarsi breves ac crassi, plantis crassis; metatarsus posticus brevis, articulis 3 sequentibus vix æquilongus, articulo 2^o brevi, subtus basi haud carinato. Abdomen vix dilatatum. Lamina supra-analis cornea, rotundato-trapezina, minute trigonali-incisa. Cerci conici, testacei. Ultimæ segmentum ventrale subconvexum, inter cercos rotundatum.

Long. 14·5; pronot. 4, latit. pronot. 5 millim.

Hab. PANAMA (*Mus. Genavense*).

2. *Calolampra bispinosa*, Sauss. (Tab. III. fig. 26.)

Calolampra bispinosa, Sauss. Societas Entomologica, 1893, p. 58.

♀. Gracilis, rufo-testacea. Caput prominulum, crassiuscule punctatum, ocellis, ore et antennis testaceis. Pronotum corneum, nitidum, fusco-castaneum, crasse punctatum, lateribus deflexis, marginibus lateralibus

* In this species the anterior femora, ♂, are armed as in *Epilampra*. The ♀ has also occasionally one spine on the posterior margin of the anterior femora. It makes a transition to the true *Epilampra*.