

testaceis, margine posteriore parum arcuato, valde obtusangulo vel rotundato. Elytra angusta, abdomen paulum superantia, testaceo-hyalina, macula basali grandi campi analis rufo-castanea. Alæ subhyalinæ, venis testaceis, margine medio costali albicante; vena ulnari ramos 3 ad marginem apicalem emittente. Pedes testacei, crassiusculi. Femora subtus spinis apicalibus minimis; femora anteriora subtus margine antico dense piloso-pectinato; femora intermedia et posteriora margine inferiore postico ante medium spinis 2 remotis armato. Tarsi sat breves, metatarso postico reliquis articulis computatis paulo brevior. Abdomen infuscatum, basi pallidum, marginibus lateralibus luteis. Lamina supra-analis testacea, transversa, margine posteriore transverse arcuato, laminæ infra-genitali æquilonga; hæc margine posteriore obtusangulo, luteo-limbato, angulo rotundato. Cerci acutiusculi, laminam supra-analem superantes. Long. 21; elytr. 18.5, pronot. 5.25, latit. pronot. 6.8 millim.

*Hab.* PANAMA (*Mus. Genavense*).

This species makes a sort of transition to the *Panchlorinæ*, the femora being very sparsely spinose, and the pronotum being intermediate in form between that of *Panchlora* and that of *Nauphoëta*.

### [OPISTHOPLATIA, Brunner de W.]

*Opisthoplatia*, Brunner de W., Syst. des Blatt. p. 198.

#### 1. *Opisthoplatia orientalis*, Burm.

*Polyzosteria orientalis*, Burm. Handb. ii. p. 482. 1; Saussure, Mém. Mex., Blatt. p. 54. 3; Stoll, Kakkerl. t. 5 d. fig. 25.

*Opisthoplatia orientalis*, Brunn. de W., Syst. Blatt. p. 199. 1; Saussure, Miss. Scient. Mex., Orthopt. p. 85. 1.

*Hab.* INDIA, in America transducta.]

### RHICNODA, Brunn. de W.

*Rhcnoda*, Brunner de W., Rev. du Syst. des Orthopt. p. 30 (1893).

A very peculiar genus, remarkable on account of the shape of the body and its produced triangular supra-anal plate.

#### *Synopsis specierum.*

*a.* Pronoti margo anterior reflexus. Femora anteriora spinis 3-4 armata, reliqua subtus inermia. Arolia inter ungues perdistincta.—1. *reflexa*, sp. n.

*aa.* Pronoti margo anterior haud vel vix reflexus.

*b.* Femora omnia subtus spinosa. Arolia inter ungues nulla (Asiaticæ).—*rugosa*, spinulosa, Br.

*bb.* Femora? Arolia? (Americanae).—[2. *laminata*, Br.]

#### 1. *Rhcnoda reflexa*, sp. n. (Tab. IV. fig. 35.)

♂. Brunneo-ochracea, subtus ochraceo-testacea; tota granulata, aptera, deplanata. Antennæ breves. Thorax valde carinatus. Pronotum rotundato-trigonale, utrinque granulatum, posterius tuberculis majoribus nitidis instructo; margine anteriore valde reflexo, disco convexo, subtrilobato. Meso- et metanotum squamoso-granulosa vel punctata, margine posteriore tuberculoso; angulis retro productis, apice rotundatis.