

rotundato; sulco anali nullo, tantum in $\frac{2}{3}$ marginis suturalis obsolete perspicuo, subperpendiculari. Alæ minutæ, ovatæ, apice late rotundatæ, campo antico quam postico valde latiore, castaneo-opaco, margine anteriore valde arcuato, campo marginali lato, reticulato. Vena discoidalis crassa, lata; vena media ad costam crasse ramosa. Vena dividens recta. Campus posterior angustus, pallidior, ab anteriore per vittam subhyalinam separatus, venis crassis, ramosis ac reticulatus; margine postico subrecto. Pedes nigro-castanei, spinis tibiis et tarsis rufescentibus. Lamina supra-analis longiuscula, rugulosa, apice incisa, marginibus lateralibus integris, angulis rotundatis. Cerci mediocres, apice rufescente.

Long. 23; elytr. 7.5, pronot. 6.2, latit. pronot. 8.4 millim.

Hab. PANAMA, Isthmus of Panama (*E. de Ville; Mus. Genavense et Mus. Bruxel-lense*).

DORYLÆA, Stål.

Dorylæa, Stål, Orthopt. Nova ex Ins. Philippinis (Öfv. k. Vetensk.-Ak. Förhandl. 1877, no. 10), p. 36.

1. *Dorylæa rhombifolia*, Stoll.

Blatta rhombifolia, Stoll, Kakkerl. t. 3 d. fig. 13; Sauss. Mém. Mex., Blatt. p. 267.

Periplaneta rhombifolia, Sauss. Miss. Scient. Mex., Orthopt. p. 72. 1.

Periplaneta histrio, Sauss. Mém. Mex., Blatt. p. 73. 18.

Periplaneta decorata, Brunn. de W. Syst. Blatt. p. 224. 2.

Hab. MEXICO; CENTRAL AMERICA.—BRAZIL.—EAST INDIES.—AFRICA.

A cosmopolitan species within the tropics.

STYLOPYGA, L. H. Fischer et auctt.

1. *Stylopyga orientalis*, Linn. et auctt.

—, Th. Moufet, Insect. sive minimorum animal. Theatrum, p. 138, figg. ♂ ♀ (1634).

Hab. NORTH AMERICA.—MEXICO; CENTRAL AMERICA.—ANTILLES.

A cosmopolitan species, inhabiting store- and dwelling-houses, ships, &c.

PERIPLANETA, Burm.

Kakerlac, Latreille; Serville.

Periplaneta, Burmeister (ex parte); Stål.

The species of this genus inhabit houses and are quite cosmopolitan; they are spread over the tropical and subtropical regions of the whole world, and are a plague in store-houses, ships, &c.

Synopsis specierum.

a. *Feminæ*. Lamina supra-analis profunde trigonali-incisa:

b. ultra segmentum ultimum ventrale producta, trigonali-bilobata.

c. Elytra tota ferruginea. Pronotum margine postico magis arcuato, rufo- vel fusco-bimaculatum vel disco rufo-fusco, fascia intramarginali circumcirca obsolete flava.—

1. *americana*, L.

BIOL. CENTR.-AMER., Orthopt., November 1893.