

- tinuous, rather broad, dusky bands; antennæ filiform; hind tibiæ furnished with about 20-23 spines in the outer row.
- b*¹. Larger; the mottling in the disc of the tegmina usually running together; antennæ slender throughout, rather long; hind tibiæ infuscated apically 1. *admirabilis*, Uhl.; *leucocera*, Stål.
- b*². Smaller; the mottlings in the disc of the tegmina appearing as distinct blotches; antennæ quite short, semiclavate; hind tibiæ unicolorous, pale 2. *pacifica*, sp. n.
- A*². Pronotum with the lateral carinæ strongly bowed inward, the dorsal field above without continuous dusky bands; antennæ with the basal joints more or less strongly flattened; hind tibiæ furnished with from 16-18 spines in the outer row.
- b*¹. Tegmina nearly or quite without mottling in the discal field.
- c*¹. Larger species. Disc of pronotum widest behind, and with lateral dusky bands showing only on the posterior lobe. Tegmina with the costal edge membranous 3. *montezuma*, Sauss.
- c*². Smaller species. Disc of pronotum as wide in front as behind, the lateral dusky bands showing on the front edge of the anterior lobe as well as on the posterior one. Tegmina with the costal edge coriaceous 4. *valida*, Rehn.
- b*². Tegmina provided with well-defined mottling in the discal field.
- c*¹. Antennæ with the basal joints only slightly depressed or flattened. Tegmina semimembranous throughout [6. *fuscovittata*, Thom.]
- c*². Antennæ with the basal joints considerably depressed or flattened, thus giving these segments a dagger-like form.
- d*¹. Large and robust; the head large and wide; pronotum with faint indications of the lateral dusky discal stripe on the sides of the extreme front edge. Wings markedly tessellate with hyaline 7. *robusta*, sp. n.
- d*². Form more slender; the head not unusually large or broad; pronotum in front without traces of lateral discal stripe.
- e*¹. Larger. The tegmina deep grass-green, rather heavily mottled with dark brown; the tibiæ and tarsi of all the legs infuscated or purplish [8. *acuticornis*, Bruner.]
- e*². Smaller. The tegmina light green, rather obscurely mottled with smoky brown; the tibiæ but little infuscated [9. *modesta*, sp. n.]

MALES.

- A*¹. Lateral carinæ of pronotum only gently bowed. Tegmina always plainly and decidedly mottled in the discal area. Wings pale fuliginous. Sides of face and thorax varied with streaks of light testaceous or dirty white.
- b*¹. Hind femora with the outer face of basal half crossed by well-