

- defined, dusky, oblique bands, the tibiae also decidedly infuscated near the middle and at the apex 1. *admirabilis*, Uhl.
- b*². Hind femora with the outer face of the basal half uniformly dusky, the tibiae less decidedly infuscated *leucocera*, Stål (?).
- A*². Lateral carinae of pronotum more decidedly bowed. Tegmina showing no decided mottling in the discal area. Wings dark fuliginous, in most cases nearly black. Sides of face and thorax not greatly varied with dirty white or testaceous.
- b*¹. Larger. Legs more or less green.
- c*¹. Hind femora on the outer face not dusky; disc in outer half of tegmina occasionally occupied by faint dusky maculations. [8. *acuticornis*, Bruner.]
- c*². Hind femora with the outer face on basal half occupied by dusky.
- d*¹. Apical third of hind femora deep transparent green, the knees not much infuscated [6. *fuscovittata*, Thom.]
- d*². Apical third of hind femora dull dirty green 3. *montezuma*, Sauss.
- b*². Smaller. Legs either infuscated or uniformly light-coloured.
- c*¹. Form robust: general colour black, varied with green on head and thorax 5. *eslavæ*, Rhen.
- c*². Form slender: general colour greenish, varied with testaceous and ferruginous [9. *modesta*, sp. n.]

1. *Syrbula admirabilis*, Uhler.

Stenobothrus admirabilis, Uhler, Proc. Ent. Soc. Phil. ii. p. 553 (1864)¹; Walk. Cat. Dermapt. Salt. Brit. Mus. iv. p. 753 (1870)²; Glover, Illustr. N. Amer. Ent., Orthopt. t. 4. fig. 13 (1872)³.

Syrbula admirabilis, Thomas, Rep. Ent. Illinois, ix. pp. 88, 93, 100–102 (1880)⁴; Bruner, Bull. Washb. Lab. Nat. Hist. i. p. 131 (1885)⁵; McNeill, Proc. Dav. Acad. Nat. Sci. vi. p. 222 (1879)⁶.

? *Syrbula leucocera*, Stål, Recens. Orthopt. i. pp. 102, 103 (1873)⁷; Thomas, Rep. U.S. Geogr. & Geol. Surv. W. 100th Merid. v. Zool. p. 869 (1875)⁸.

Hab. NORTH AMERICA^{1–6}, United States, chiefly southward.—MEXICO, San Rafael in Vera Cruz (*coll. Bruner*).

In the collection of the writer there are several males from San Rafael, Vera Cruz, which differ but little from others taken in Florida. Perhaps the *S. leucocera* of Stål may be identical with these Mexican specimens. No females have been examined from Mexico.

2. *Syrbula pacifica*, sp. n.

General colour grass-green; the pronotum with the lateral carinae nearly straight, its disc laterally marked with longitudinal dark bands. Tegmina provided with a row of large circular dusky spots along the discal field, occasionally running together as in *S. admirabilis* and several of the other species. Wings only faintly clouded or fuscous. Head of moderate size; the occiput short, about two-thirds as long as