

2. *Conalcæa truncatipennis*, Scudd.

Conalcæa truncatipennis, Scudd. Proc. U.S. Nat. Mus. xx. p. 25 (1897)¹.

Hab. MEXICO, Saltillo, Coahuila¹.

Not represented in the material studied.

3. *Conalcæa neomexicana*, Scudd.

Conalcæa neomexicana, Scudd. Proc. U.S. Nat. Mus. xx. p. 26, t. 2. fig. 9 (1897)¹; Caudell, Proc. U.S. Nat. Mus. xxviii. p. 476 (1905)².

? *Pezotettix humphreysii*, Thom. Rep. Geogr. and Geol. Expl. W. 100th Merid. v. p. 890, t. 45. fig. 1 (1875)³ (in part.).

Hab. NORTH AMERICA, Silver City in New Mexico¹, Huachuca Mts.² and Phoenix in Arizona (*coll. L. Bruner*).

This insect, like *C. miguelitana*, is very variable in size and colour. While the type only measured 19, the Phoenix specimen quoted is 25 millimetres in length. *C. neomexicana* will certainly be found in the adjacent portion of Mexico.

[BARYTETTIX, Scudder.

Barytettix, Scudd. Proc. Amer. Acad. Art. Sci. xxxii. no. 9, p. 197 (Jan. 1897); Proc. U.S. Nat. Mus. xx. pp. 10, 27 (1897).

The members of the genus *Barytettix* remind one a little of a small *Brachystola* in their general appearance. The two known species appear to be confined to the Pacific coast region in the vicinity of Cape St. Lucas.

1. *Barytettix crassus*, Scudd. (Tab. IV. figg. 8, 8 *a*, *b*, ♂; 9, ♀.)

Barytettix crassus, Scudd. Proc. U.S. Nat. Mus. xx. p. 28, t. 2. fig. 10 (1897)¹.

Hab. LOWER CALIFORNIA, San José del Cabo¹.

There are a number of specimens of both sexes in the present writer's collection. The females are much larger and more robust than the opposite sex, some of them measuring fully 40 millimetres in length.

2. *Barytettix peninsulæ*, Scudd.

Barytettix peninsulæ, Scudd. Proc. U.S. Nat. Mus. xx. p. 28 (1897)¹.

Hab. LOWER CALIFORNIA¹.

Only the type is at hand. It is much smaller than the preceding species.]