

10. **Melanoplus bowditchi**, Scudd.

*Melanoplus bowditchi*, Scudd. Proc. Bost. Soc. Nat. Hist. xx. p. 72 (1879)<sup>1</sup>; Proc. U.S. Nat. Mus. xx. p. 157, t. 11. fig. 3 (1897)<sup>2</sup>.

*Hab.* NORTH AMERICA, Texas, New Mexico, &c. (*coll. L. Bruner*).—MEXICO, Northern Chihuahua (*L. Bruner*).

11. **Melanoplus flavidus**, Scudd.

*Melanoplus flavidus*, Scudd. Proc. Bost. Soc. Nat. Hist. xx. p. 74 (1879)<sup>1</sup>; Proc. U.S. Nat. Mus. xx. p. 158, t. 11. fig. 4 (1897)<sup>2</sup>; Rehn, Proc. Acad. Nat. Sci. Philad. 1907, p. 54<sup>3</sup>.

*Hab.* NORTH AMERICA, various localities from Nebraska to Southern Texas and Arizona<sup>3</sup> (*coll. L. Bruner*).—MEXICO, Tampico, Tamaulipas (*coll. Ill. State Lab. Nat. Hist.*).

12. **Melanoplus elongatus**, Scudd.

*Melanoplus elongatus*, Scudd. Proc. U.S. Nat. Mus. xx. p. 160, t. 11. fig. 5 (1897)<sup>1</sup>.

*Hab.* NORTH AMERICA, Kansas and New Mexico<sup>1</sup>.—MEXICO, Villa Lerdo, Durango (*L. Bruner*), Bledos, San Luis Potosi<sup>1</sup> (*Dr. Palmer*), Monterey, Nuevo Leon (*Rehn*), Guanajuato (*A. Dugès*).

13. **Melanoplus glaucipes**, Scudd.

*Caloptenus glaucipes*, Scudd. Proc. Bost. Soc. Nat. Hist. xvii. p. 476 (1875)<sup>1</sup>; Ent. Notes, iv. p. 75 (1875)<sup>2</sup>; Thom. Rep. U.S. Ent. Comm. i. p. 42 (1878)<sup>3</sup>.

*Melanoplus glaucipes*, Scudd. Canad. Ent. xii. p. 75 (1880)<sup>4</sup>; Proc. U.S. Nat. Mus. xx. p. 161, t. 11. fig. 6 (1897)<sup>5</sup>.

*Hab.* NORTH AMERICA<sup>1-5</sup>.—MEXICO, Villa Lerdo, Durango (*L. Bruner*).

14. **Melanoplus atlanis**, Riley.

*Caloptenus atlanis*, Riley, Ann. Rep. St. Ent. Mo. vii. p. 169 (1875)<sup>1</sup>; Thom. Bull. Ill. Mus. Nat. Hist. i. p. 68 (1876)<sup>2</sup>.

*Melanoplus atlanis*, Scudd. Rep. U.S. Ent. Comm. ii. Append. p. 24, t. 17. fig. 6 (1881)<sup>3</sup>; Proc. U.S. Nat. Mus. xx. p. 178, t. 12. fig. 7 (1897)<sup>4</sup>.

*Hab.* NORTH AMERICA, various localities from the Atlantic to the Pacific and from the Saskatchewan to the Rio Grande<sup>1-4</sup>.—MEXICO, numerous places between the Rio Grande and Orizaba, Amula, and Omilteme, Guerrero (*H. H. Smith*).

This is one of the most widely distributed members of the genus, and is also variable in size and colour. The synonymy is much more completely worked out in Scudder's Monograph.

15. **Melanoplus spretus**, Uhler.

*Caloptenus spretus*, Uhler, MS. (1863).

*Acridium spretis*, Thom. Trans. Ill. St. Agr. Soc. v. p. 450 (1865)<sup>1</sup>.