

*Hab.* NORTH AMERICA, from the Atlantic to the Pacific and from the Saskatchewan to the Rio Grande <sup>4</sup>.—MEXICO, Queretaro, Guanajuato, Atoyac, Vera Cruz (*coll. L. Bruner*).

This insect, along with *M. atlanis* and *M. bivittatus*, enjoys the most extended distribution of all the species of the genus. It has been the most frequently mentioned in entomological literature, and has a very extended synonymy, which may be found in Scudder's paper <sup>4</sup> cited above.

### 35. *Melanoplus terminalis*, Scudd.

*Melanoplus terminalis*, Scudd. Proc. U.S. Nat. Mus. xx. p. 293, t. 19. fig. 7 (1897) <sup>1</sup>.

*Hab.* NORTH AMERICA, Gulf coast of Texas <sup>1</sup>, extreme South-western Texas (*coll. L. Bruner*).

While no records or specimens of this insect are at hand from Mexican territory, it most assuredly occurs across the Rio Grande in that country.

### 36. *Melanoplus cyanipes*, Scudd.

*Melanoplus cyanipes*, Bruner, MS. <sup>1</sup>.

*Melanoplus cyanipes*, Scudd. Proc. U.S. Nat. Mus. xx. p. 295, t. 19. fig. 8 (1897) <sup>2</sup>.

*Hab.* NORTH AMERICA, Los Angeles and San Diego, California <sup>2</sup> (*coll. L. Bruner*)  
LOWER CALIFORNIA, San José del Cabo.—MEXICO, Tepic, Jalisco (*colls. L. Bruner and Calif. Acad. Nat. Sci.*).

### 37. *Melanoplus cinereus*, Scudd.

*Melanoplus cinereus*, Scudd. Proc. Bost. Soc. Nat. Hist. xix. pp. 288, 290 (1878) <sup>1</sup>; Proc. U.S. Nat. Mus. xx. p. 296, t. 19. fig. 9 (1897) <sup>2</sup>.

*Caloptenus cinereus*, Riley, in Kingsley's Stand. Nat. Hist. ii. p. 195 (1884) <sup>3</sup>.

*Hab.* NORTH AMERICA, numerous localities between the 49th degree of latitude and the Mexican boundary <sup>2</sup> (*coll. L. Bruner*).

Since this insect is distributed over the Rocky Mountain region wherever "sagebrush" grows, it most certainly extends along the tablelands into Northern Mexico.

### 38. *Melanoplus complanatipes*, Scudd.

*Melanoplus complanatipes*, Scudd. Proc. U.S. Nat. Mus. xx. p. 298, t. 19. fig. 10 (1897) <sup>1</sup>.

*Hab.* LOWER CALIFORNIA <sup>1</sup>, Cape St. Lucas <sup>1</sup>.—MEXICO, Sonora <sup>1</sup>.

### 39. *Melanoplus impiger*, Scudd.

*Melanoplus impiger*, Scudd. Proc. U.S. Nat. Mus. xx. p. 306, t. 20. figg. 7, 8 (1897) <sup>1</sup>.

*Hab.* NORTH AMERICA, Gulf Coast and many other Texas localities <sup>1</sup> (*coll. L. Bruner*).

Since this insect occurs in extreme Southern Texas on the Rio Grande, it must be also present in the adjacent Mexican territory.