

40. **Melanoplus corpulentus**, Scudd.

Melanoplus corpulentus, Bruner, MS.¹.

Melanoplus corpulentus, Scudd. Proc. U.S. Nat. Mus. xx. p. 313, t. 20. fig. 10 (1897)²; Rehn, Proc. Acad. Nat. Sci. Philad. 1907, p. 55³.

Hab. NORTH AMERICA, Silver City² and Las Vegas, New Mexico (*Cockerell*), Huachuca Mts., Arizona³. — MEXICO, Sonora, Zacatecas, Tlalpam, &c.², Guadalajara, Jalisco³ (*Rehn*).

41. **Melanoplus variabilis**, Scudd.

Pezotettix variabilis, Bruner, MS.¹.

Melanoplus variabilis, Scudd. Proc. U.S. Nat. Mus. xx. p. 319, t. 21. fig. 8 (1897)².

Hab. MEXICO, city of Mexico², Queretaro², &c. (*coll. L. Bruner*).

This insect is very common in the regions where it occurs, and is very variable in colour, hence the specific name.

42. **Melanoplus plebejus**, Stål.

Pezotettix plebejus, Stål, Bihang till K. Svensk. Vet.-Ak. Handl. v. no. 9, p. 12 (1878)¹.

Pezotettix pupæformis, Scudd. Proc. Bost. Soc. Nat. Hist. xx. pp. 83, 84 (1879)².

Melanoplus plebejus, Scudd. Proc. U.S. Nat. Mus. xx. p. 326, t. 22. fig. 2 (1897)³.

Hab. NORTH AMERICA, Texas¹⁻³. — MEXICO, Victoria, Tamaulipas (*coll. Ill. St. Lab. Nat. Hist.*).

43. **Melanoplus marginatus**, Scudd.

Pezotettix marginatus, Scudd. Ann. Rep. Chief Eng. 1876, p. 504 (1876)¹.

Melanoplus marginatus, Scudd. Proc. U.S. Nat. Mus. xx. p. 330, t. 1. fig. *i* and t. 22. fig. 5 (1897)².

Hab. NORTH AMERICA, various places in Southern California^{1 2}, Los Angeles and San Diego (*coll. L. Bruner*).

This insect probably extends into Mexican territory.

44. **Melanoplus arizonæ**, Scudd.

Melanoplus arizonæ, Scudd. Proc. Bost. Soc. Nat. Hist. xx. pp. 64, 65 (1879)¹; Proc. U.S. Nat. Mus. xx. p. 340 (1897)²; Rehn, Trans. Amer. Ent. Soc. xxvii. p. 228 (1901)³.

Hab. NORTH AMERICA, Arizona^{1 2}. — MEXICO, Cuernavaca, Morelos³.

The Mexican citation may be an error, though this is only a surmise on the part of the present writer.

45. **Melanoplus differentialis**, Uhler.

Caloptenus differentialis, Uhler, MS. (1863)¹.

Acridium differentiale, Thom. Trans. Ill. St. Agric. Soc. v. p. 450 (1865)².

Cyrtacanthacris differentialis, Walk. Cat. Dermapt. Salt. Brit. Mus. iv. p. 610 (1870)³.