

2. **Libethra confusa**, Br.*Libethra confusa*, Brunner, op. cit. ii. p. 308 (1907).*Hab.* GUATEMALA (*Mus. Vienna*; *Mus. Geneva*).3. **Libethra elegantior**, Br.*Libethra elegantior*, Brunner, op. cit. ii. p. 308 (1907).*Hab.* MEXICO, Omilteme in Guerrero 8000 feet (*H. H. Smith, in coll. Godman & Salvin*: 1 ♀, type).4. **Libethra regularis**, Br.*Libethra regularis*, Brunner, op. cit. ii. p. 308 (1907).*Hab.* MEXICO (*Mus. Budapest*: ♀).—TRINIDAD, Port of Spain (*Mus. Hamburg*: ♀ type).5. **Libethra brevipes**, Br.*Libethra brevipes*, Brunner, op. cit. ii. p. 309 (1907).*Hab.* MEXICO (*Mus. Paris*).

## OCNOPHILA, Br.

*Ocnophila*, Brunner, op. cit. ii. p. 309 (1907).*Conspectus specierum.*

1. Caput haud lobatum, interdum spinosum.
2. Femora intermedia vel postica superne lobis singulis ornata.
3. Caput inter oculos bispinosum. Mesonotum et segmenta abdominalia integra . . . . . *O. crudis*, Br.
- 3'. Caput in occipite spinosum. Mesonotum et segmenta abdominalia lobata.
4. Picea. Occiput 4-spinosum. Pronotum rugosum . . . . . *O. strumosa*, Br.
- 4'. Albo-pruinosa. Occiput multispinosum. Pronotum biserialiter dense spinulosum . . . . . *O. ornatissima*, Br.
- 2'. Femora omnia superne integra vel lobulis minimis regulariter dispositis instructa, interdum subtus lobulata vel lobulato-dentata.
3. Vertex muticus, interdum granulis majoribus obsitus.
4. Caput granulis majoribus obsitum, haud carinatum.
5. Abdomen totum distincte multicarinatum. Femora submutica. [*O. submutica*, Br.]
- 5'. Abdomen indistincte carinatum. Femora intermedia et postica subtus minime lobulata . . . . . *O. medians*, Br.
- 4'. Caput minime granulatum vel læve, carinatum . . . . . *O. ciliata*, Br.
- 3'. Vertex bispinosus . . . . . *O. godmani*, Br.
- 1'. Caput in occipite bi-auriculatum . . . . . *O. mexicana*, Sauss.
- Species incertæ sedis . . . . . *O. aurita*, Rehn.