

1. **Dyme grossetuberculata**, Br.

Dyme grossetuberculata, Brunner, op. cit. ii. p. 323 (1907).

Hab. HONDURAS (*Mus. Vienna*).

2. **Dyme discors**, Br.

Dyme discors, Brunner, op. cit. ii. p. 324 (1907).

Hab. PANAMA, Volcan de Chiriqui (*Champion, in coll. Godman & Salvin*: 5 ♂ ♂).

3. **Dyme modesta**, Br.

Dyme modesta, Brunner, op. cit. ii. p. 324 (1907).

Hab. PANAMA, Chiriqui (*Champion, in coll. Godman & Salvin*: 1 ♂, type).

4. **Dyme chiriquensis**, Br.

Dyme chiriquensis, Brunner, op. cit. ii. p. 325 (1907).

♀. Lævis, colore nitido, olivaceo. Antennæ longiores quam pedibus anticis. Femora intermedia et (?) postica mutica. Abdomen haud carinatum. Segmentum anale carinatum, apice emarginatum, lamina supra-analis triangularis, minuta. Operculum haud carinatum, lanceolatum, apice obtuso, apicem abdominis attingens. Cerci breves, teretes, subacuminati.

Long. corp. 100.1 mm.

„ meson. 25 mm.

„ metan. c. segm. med. 17 mm.

„ fem. ant. 26 mm.

Hab. GUATEMALA, San Gerónimo (*Champion, in coll. Godman & Salvin*: ♀); PANAMA, Chiriqui (*Mus. Vienna*: ♂ type).

The description of the female is drawn up from a single example in the Godman and Salvin collection labelled “*Dyme lævis*” by Brunner; no species of this name appears in his Monograph, and the specimen cannot be referred to any of the forms described therein from females only. I am almost certain that this is the female of *D. chiriquensis*, Br., known only by the male, and as such have described it.

5. **Dyme incolumis**, Br.

Dyme incolumis, Brunner, op. cit. ii. p. 326 (1907).

Hab. GUATEMALA, Panzos in Vera Paz (*Conradt, in coll. Godman & Salvin*: 1 ♂, type).

6. **Dyme depressa**, Br.

Dyme depressa, Brunner, op. cit. ii. p. 327 (1907).

Hab. GUATEMALA, Las Mercedes, Cerro Zunil, Purula in Vera Paz (*Champion, in coll. Godman & Salvin*: 3 ♀ ♀, including type).