

**7. *Bacunculus carinulatus*, Br.**

*Bacunculus carinulatus*, Brunner, op. cit. ii. p. 334 (1907).

*Hab.* MEXICO, Ciudad in Durango 8100 feet (*Forrer*), Sierra de las Aguas Escondidas in Guerrero 7000 feet (*H. H. Smith*) [*coll. Godman & Salvin*: 2 ♂♂, types].

**8. *Bacunculus godmani*, Br.**

*Bacunculus godmani*, Brunner, op. cit. ii. p. 334 (1907).

*Hab.* MEXICO, Amula in Guerrero 6000 feet (*H. H. Smith*), Jalisco (*Schumann*) [*coll. Godman & Salvin*: 3 ♂♂ (types), 2 larvæ, ♂ & ♀].

**9. *Bacunculus tridens*, Burm.**

*Bacteria tridens*, Burmeister, Handb. Ent. ii. p. 567 (1839).

*Bacunculus tridens*, Brunner, op. cit. ii. p. 334 (1907).

*Hab.* MEXICO (*Mus. Berlin, Mus. Brit., Mus. Paris, Geneva, Vienna*), Amula, Omilteme, and Chilpancingo in Guerrero 4600–8000 feet (*H. H. Smith*) [*coll. Godman & Salvin*: 4 ♂♂, 1 ♀].

**10. *Bacunculus elongatus*, Br.**

*Bacunculus elongatus*, Brunner, op. cit. ii. p. 335 (1907).

*Hab.* MEXICO, Dos Arroyos in Guerrero 1000 feet (*H. H. Smith, in coll. Godman & Salvin*: 1 ♂, type).

**11. *Bacunculus contractus*, Br.**

*Bacunculus contractus*, Brunner, op. cit. ii. p. 335 (1907).

*Hab.* MEXICO, Sierra de Nayarit in Jalisco (*Mus. Paris*).

**12. *Bacunculus ignavus*, Rehn.**

*Heteronemia ignava*, Rehn, Proc. Acad. Nat. Sci. Philad. 1904, p. 54.

*Hab.* COSTA RICA, Piedras Negras (*U.S. Nat. Mus.*).

**13. *Bacunculus olmecus*, Sauss.**

*Bacteria olmecus*, Saussure, Miss. Sci. Mex., Orthopt. p. 156 (1870).

*Bacunculus olmecus*, Brunner, op. cit. ii. p. 335 (1907).

*Hab.* MEXICO, Eastern Cordillera (*Mus. Paris*).

**14. *Bacunculus forcipatus*, Br.**

*Bacunculus forcipatus*, Brunner, op. cit. ii. p. 335 (1907).

*Hab.* MEXICO, San Salvador (*Mus. Vienna*).

**15. *Bacunculus incongruens*, Br.**

*Bacunculus incongruens*, Brunner, op. cit. ii. p. 336 (1907).