

6. **Diapheromera beckeri**, Kaup.*Diapheromera beckeri*, Kaup, Berl. ent. Zeitschr. xv. p. 27 (1871).*Hab.* MEXICO.7. **Diapheromera (?) bidens**, Kaup.*Diapheromera bidens*, Kaup, t. c. p. 28 (1871).*Hab.* MEXICO.8. **Diapheromera (?) strigiceps**, Kaup.*Diapheromera strigiceps*, Kaup, t. c. p. 28.*Hab.* MEXICO.Subfam. *PHIBALOSOMINÆ*.*Conspectus generum.*

1. Femora antica superne haud compressa vel superne utrinque sub-æqualiter spinosa. (Sect. EURYCANTHÆ.)
2. Femora antica haud oblique compressa, superne lobato-dentata . . . PERICENTRUS, Redt.
- 2'. Femora antica oblique compressa.
3. Occiput bicornè.
4. Apex abdominis ♀ rostratus RHYNCHACRIS, Redt.
- 4'. Apex abdominis ♀ haud rostratus PHANTASIS, Sauss.
- 3'. Occiput spinosum vel tuberculatum, haud bicornè.
4. Mesonotum strumosum. Alæ ♂ ♀ distinctæ, lobiformes . . . HYPOCYRTUS, Redt.
- 4'. Mesonotum haud strumosum. Alæ ♂ ♀ desunt [LAMPONIUS, Stål.]
- 1'. Femora antica superne plus minusve compressa rite inermia. (Sect. PHIBALOSOMATA.)
2. Femora antica carina infera mediana margini antico appropinquata; femora 4 postica subtus ante apicem in carina externa dentibus vel spinis majoribus haud instructa. ♀ aptera.
3. Metatarsus superne sulcatus CLONISTRIA, Stål.
- 3'. Metatarsus superne carinatus vel cristatus.
4. Segmentum medianum metanoto distincte brevius. ♂ apterus. BOSTRA, Stål.
- 4'. Segmentum medianum metanoto æquilongum vel longius. ♂ plerumque alatus BACTERIA, Latr.
- 2'. Femora antica carina infera mediana in medio sita; femora 4 postica margine laterali infero saltem externo ante apicem dentibus vel spinis majoribus 1-2 armato. ♂ ♀ alatus.
3. Femora omnia foliaceo-lobata, antica cum tibiis undulato-dilatata . . . PTERINOXYLUS, Serv.
- 3'. Femora antica cum tibiis haud foliaceo-lobata, 4 postica spinosa . . . HAPLOPUS, Gray.