

5. **Thalictrum longistylum**, DC. Syst. i. p. 171; Deless. Ic. Sel. i. t. 7?

SOUTH MEXICO, Chiapas (*Ghiesbreght*), Oaxaca (*Ghiesbreght*), Orizaba (*Bourgeau*, 2726),—and southward to PERU. Hb. Kew.

6. **Thalictrum mexicanum**, DC. Syst. i. p. 187.

MEXICO (*Hernandez*).

7. **Thalictrum peltatum**, DC. Prod. i. p. 11.

MEXICO (*De Bergher*, 3).

8. **Thalictrum pubigerum**, Benth. Pl. Hartw. p. 285.

SOUTH MEXICO, on the banks of rivers, Lagos (*Hartweg*, 1591). Hb. Kew.

9. **Thalictrum rutidocarpum**, DC. Syst. i. p. 172.

SOUTH MEXICO (*Sallé, Graham*), Orizaba (*Botteri*, 824).—Mountains of TROPICAL SOUTH AMERICA. Hb. Kew.

10. **Thalictrum strigillosum**, Hemsley, Diag. Pl. Nov. pars 1, p. 1.

Caule ramisque plus minusve strigilloso, foliolis graciliter petiolulatis subtus hispidis supra sparse puberulis, floribus longe pedicellatis, antheris insigniter aristatis, stylo pubescente longissime exserto, achæniis fere sessilibus rugis elevatis reticulatis notatis.

*Herba* 2–3-pedalis, ramosa, strigilloso-hispida. *Radix* subcarnosa, fibrosa. *Caulis* sulcato-striatus, strigillosus vel demum fere glaber. *Folia* (radicalia non vidi) petiolata, subquadri-ternatim divisa, foliolis petiolulatis, cordato-rotundatis vel basi cuneatis, apice 3–5-lobatis, ad 8–12 lineas latis, præcipue subtus secus nervos hispidis. *Flores* longe pedicellati, pedicellis basi bracteis parvis munitis; sepala glabra, ovato-oblonga, ad 3 lin. longa; stamina calycem superantia, filamentis capillaceis, antheris aristatis filamentis æquilongis. *Achenia* circiter 7, sessilia vel brevissime stipitata, oblique lunata, 2–3 lin. longa, insigniter costato-reticulata.

SOUTH MEXICO, rare in ravines among bushes (*Schaffner*), Tizapan, valley of Mexico (*Bourgeau*, 276), Zimapan (*Coulter*, 652), mountains around Mitla, Oaxaca (*Andrieux*, 546), between San Miguel and La Joya (*Schiede*). Hb. Kew.

This differs from its allies *T. rutidocarpum*, *longistylum*, &c. in being more or less clothed with hispid hairs, which on the stem are long and on the leaves short, and in the leaflets, anthers, and achenes. Perhaps it will eventually be found necessary to unite several of these forms as one variable species.

11. **Thalictrum wrightii**, A. Gray, Pl. Wright. ii. p. 7.

NORTH MEXICO, mountain ravine at Santa Cruz, Sonora (*Wright*). Hb. Kew.

12. **Thalictrum**, sp.

SOUTH MEXICO, between Chalco and Gonacatepec (*Andrieux*, 545). Hb. Kew.

13. **Thalictrum**, sp.

SOUTH MEXICO, between San Miguel and La Joya (*Schiede*). Hb. Kew.