

14. **Thalictrum**, sp.

SOUTH MEXICO, Vera Cruz (*Galeotti*, 4541 and 4549, *Linden*, 963). Hb. Kew.
 "Very near *T. peltatum*, DC., but leaflets not peltate."—*Bentham* in Hb. Kew.

15. **Thalictrum**, sp. (foliolis amplis latis).

SOUTH MEXICO, Oaxaca (*Ghiesbreght*). Hb. Paris.

3. ANEMONE.

Anemone, Linn. Gen. Plant. n. 694; Benth. et Hook. Gen. Plant. i. p. 4.

About seventy species, chiefly in the temperate and cold regions of the northern hemisphere, a few scattered in the mountainous regions of Central and South America; three occur in South Africa, and one in Australia.

1. **Anemone caroliniana**, Walt. Carol. p. 157; Marcy's Exploration of the Red River, t. 1.

Anemone tenella, Pursh.

NORTH AMERICA, from North Carolina through the south-eastern States to—NORTH MEXICO, Chihuahua (*Torrey*).

2. **Anemone mexicana**, H. B. K. Nov. Gen. et Sp. v. p. 41.

NORTH AMERICA, Illinois, Louisiana, and Texas to—SOUTH MEXICO, near Santa Rosa (*Humboldt & Bonpland*), Oaxaca (*Galeotti*), Zimapan (*Coulter*, 654), Orizaba (*Botteri*, 21), Vera Cruz (*Linden*, 964). Hb. Kew.

4. MYOSURUS.

Myosurus, Linn. Gen. Plant. n. 394; Benth. et Hook. Gen. Plant. i. p. 5.

Two species, both annual herbs. *M. minimus* is widely dispersed in the temperate parts of the northern hemisphere, and also occurs in South America and Australia. Probably introduced in some of the localities. *M. aristatus* grows in Western Extra-tropical America, both North and South, and also in New Zealand, and is likely to occur in Mexico.

1. **Myosurus minimus**, Linn. Sp. Pl. 407; Gray, Gen. Ill. t. 8.

Temperate regions of the northern hemisphere, AUSTRALIA and SOUTH AMERICA; NORTH AMERICA southward to—NORTH MEXICO, Chihuahua (*Bigelow*), Sonora (*Schott*). Hb. Kew.

Tribe RANUNCULEÆ.

5. RANUNCULUS.

Ranunculus, Linn. Gen. Plant. n. 699; Benth. et Hook. Gen. Plant. i. p. 5.

Annual and perennial herbs; about 160 species dispersed over nearly all temperate