

sesquipollicaribus, calcari basi integro leviter curvato deorsum spectante sepalis longiore, calcaribus petalinis calcari calycino æquilongis ultra medium tubulosis, petalis lateralibus exappendiculatis, carpellis 3 puberulis.

Herba biennis seu perennis, 2–3 ped. alta, caule tereti glabro lævi vix ramoso. *Folia* glabra, alte palmatim 5-partita, segmentis alte trisectis, laciniis ultimis linearibus obtusis, lamina $1\frac{1}{2}$ –3 poll. diametro, petiolo foliorum radicalium gracili 5–6-pollicari, superiorum gradatim brevior. *Pedunculi* elongati, 2–4-flori vel interdum probabiliter pluriflori. *Flores* distantes, puberuli, longe graciliterque pedicellati circiter sesquipollicares, pedicellis puberulis 2–3-pollicaribus, bracteis bracteolisque lineari-subulatis; sepala glabrescentia, oblongo-elliptica, apice rotundata, semipollicaria, calcari basi integro leviter curvato deorsum spectante sepalis longiore; petalorum superiorum laminis oblique oblongis, apice breviter bifidis fere glabris quam sepala brevioribus, calcaribus calcari calycino æquilongis, ultra medium tubulosis; petala lateralia unguiculata, basi exappendiculata, intus leviter barbata, alte bifida, lobis acutis; stamina glabra. *Carpella* 3, immatura, puberula.

NORTH MEXICO, region of San Luis Potosi, 6000 to 8000 feet (*Parry & Palmer*, 7).
Hb. Kew.

This species is characterized by the long narrow lobes of its palmately-divided leaves, by the spurs of the petals equalling the calyx-spur and tubular from the base to above the middle, and by the lateral petals being deeply bifid, not furnished with an appendage at the base, and only slightly bearded.

4. **Delphinium pedatisectum**, Hemsley, Diag. Pl. Nov. pars alt. p. 18.

Foliis caulinis pedatisectis, segmentis 7–3 integerrimis, floribus 1– $1\frac{1}{4}$ poll. longis, calcari calycino basi integro leviter curvato deorsum spectante sepalis longiore, calcaribus petalinis calcari calycino fere tertia parte brevioribus vix ad medium tubulosis, petalis lateralibus basi minute appendiculatis, carpellis 3 immaturis pubescentibus.

Herba erecta, ramosa, ramis teretibus glaberrimis lævibus. *Folia* radicalia ignota, caulina inferiora pedatisecta 7-loba, superiora trisecta, lobis linearibus 1–2 poll. longis, subtus pubescentibus. *Pedunculi* elongati, circiter 4-flori. *Flores* distantes, puberuli, longe graciliterque pedicellati, circiter pollicares, pedicellis puberulis, bracteis bracteolisque lineari-subulatis; sepala glabrescentia oblique oblonga, calcari basi integro leviter curvato deorsum spectante; petalorum superiorum laminis angustis alte bifidis, lobis acutis, calcaribus calcari calycino multo brevioribus non vel vix ad medium tubulosis extus intusque puberulis; petala lateralia unguiculata, intus barbata, limbo bipartito, ungue basi appendice squamiformi ornato; stamina glabra. *Carpella* 3, immatura, tomentosa, stylis elongatis.

MEXICO (*Parkinson*), without locality. Hb. Kew.

Distinguished by the pedatisect leaves, by the spurs of the deeply bifid upper petals being shorter than the calyx-spur, and by the bipartite lateral petals having a scale-like appendage at the base.

5. **Delphinium wislizeni**, Engelm. Bot. Wisliz. Exp. p. 22.

NORTH MEXICO, Cosiquiriachi (*Wislizenus*).