

4. **Anona echinata**, Dunal, Monogr. Anon. p. 68, t. 4.

Widely dispersed in TROPICAL SOUTH AMERICA and reaching—PANAMA, Bujio railway-station (*S. Hayes*, 142). Hb. Kew.

5. **Anona excelsa**, H. B. K. Nov. Gen. et Sp. v. p. 59.

SOUTH MEXICO, near La Venta del Exido (*Humboldt & Bonpland*).

6. **Anona globiflora**, Schl. in Linnæa, x. p. 235.

SOUTH MEXICO, Papantla (*Liebmann*), near the Hacienda de la Laguna (*Schiede*). Hb. Kew.

7. **Anona involucrata**, Baill. Adans. viii. p. 265.

SOUTH MEXICO, Tlatatla (*Liebmann*). Hb. Kew.

8. **Anona liebmänniana**, Baill. Adans. viii. p. 266.

SOUTH MEXICO, Comaltepec (*Liebmann*).

9. **Anona muricata**, Linn. Sp. Pl. p. 756; Jacq. Obs. i. t. 5.

Anona bonplandiana, H. B. K.

Native of Tropical America, and commonly cultivated. Perhaps not indigenous in CENTRAL AMERICA and MEXICO. Propanche de la Concepcion (*Liebmann*). Hb. Kew.

Grisebach thinks this may be indigenous only in some of the West-Indian Islands.

10. **Anona palustris**, Linn. Sp. Pl. p. 757.

Also commonly cultivated.

SOUTH MEXICO, Vera Cruz (*Schiede*).

11. **Anona purpurea**, DC. Prodr. i. p. 84; Dunal, Monogr. Anon. t. 2.

SOUTH MEXICO (*Moçino & Sessé*); PANAMA (*Duchassaing*). Hb. Kew.

12. **Anona reticulata**, Linn. Sp. Pl. p. 757.

PANAMA, Island of Taboga (*Seemann*). Hb. Kew.

Commonly cultivated for its fruit, and widely dispersed in Tropical America.

13. **Anona squamosa**, Linn. Sp. Pl. p. 757; Jacq. Obs. i. t. 6. fig. 1.

Cultivated. Hb. Kew.

14. **Anona**, sp.

PANAMA, Chagres (*Fendler*). Hb. Kew.

15. **Anona**, sp. (aff. *A. sylvatica*, A. St.-Hil.).

SOUTH MEXICO, region of Orizaba (*Bourgeau*, 2482). Hb. Kew.

16. **Anona**, sp.

SOUTH MEXICO (*Hahn*). Hb. Kew.

17. **Anona**, sp. (? *A. sericeæ* var. *foliis pedalibus*).

PANAMA, Obispo Falls (*S. Hayes*, 127). Hb. Kew.