

tropical America; a few occur in the Canary Islands, South Africa, and the Mascarene Islands.

1. **Ocotea effusa**, Hemsley.

*Oreodaphne effusa*, Meissn. in DC. Prodr. xv. 1, p. 120.

SOUTH MEXICO, Sierra San Pedro Nolasco &c. (*Jurgensen*, 906), Vera Cruz (*Linden*, 17), Jalapa 4000 feet (*Galeotti*, 7061). Hb. Kew.

2. **Ocotea helicterifolia**, Hemsley.

*Oreodaphne helicterifolia*, Meissn. in DC. Prodr. xv. 1, p. 123.

SOUTH MEXICO, San Bartolo, Chiapas (*Linden*, 1641). Hb. Kew.

3. **Ocotea klotzschiana**, Hemsley.

*Oreodaphne klotzschiana*, Nees in Linnæa, xxi. p. 523; DC. Prodr. xv. 1, p. 128.

*Oreodaphne psychotrioides*, Benth. Pl. Hartw. p. 50, non Nees.

SOUTH MEXICO, region of Orizaba (*Bourgeau*, 2517), Banco (*Hartweg*). Hb. Kew.

4. **Ocotea mexicana**, Hemsley.

*Oreodaphne mexicana*, Meissn. in DC. Prodr. xv. 1, p. 118.

a. **subsessilis**, Meissn. loc. cit.

SOUTH MEXICO, Cordillera of Oaxaca, 3000 feet (*Galeotti*, 7004), Sierra San Pedro Nolasco &c. (*Jurgensen*, 937). Hb. Kew.

β. ? **longipes**, Meissn. loc. cit.

SOUTH MEXICO, around Orizaba (*Botteri*, 1018). Hb. Kew.

γ. **diminuta**, Meissn. loc. cit.

SOUTH MEXICO, Sierra San Pedro Nolasco &c. (*Jurgensen*, 575). Hb. Kew.

5. **Ocotea mucosa**, Hemsley.

*Oreodaphne mucosa*, Nees, Syst. Laur. p. 401; DC. Prodr. xv. 1, p. 133.

NEW SPAIN (hb. *Pavon*).

6. **Ocotea psychotrioides**, H. B. K. Nov. Gen. et Sp. ii. p. 162.

*Oreodaphne psychotrioides*, Nees, Syst. Laur. p. 452; DC. Prodr. xv. 1, p. 124 (varietates).

SOUTH MEXICO, region of Orizaba (*Botteri*, 1040, 1041, 1042, 1043, 1050; *Sumichrast*, 940), Jalapa (*Linden*, 7023, 7047; *Humboldt & Bonpland*), Cuesta de Naulingo (*Schiede*). Hb. Kew.

7. **Ocotea sieberi**, Hemsley.

*Oreodaphne sieberi*, Meissn. in DC. Prodr. xv. 1, p. 137.

SOUTH MEXICO, Teapa, Tabasco (*Linden*, 1607; *Johnson*, 127); PANAMA, Lion Hill (*S. Hayes*, 384)?—WEST INDIES; BRAZIL. Hb. Kew.