

8. **Ocotea subtriplinervia**, Hemsley.*Oreodaphne subtriplinervia*, Meissn. in DC. Prodr. xv. 1, p. 125.SOUTH MEXICO, near Jalapa, Vera Cruz (*Linden*, 17; *Galeotti*, 7061).9. **Ocotea tampicensis**, Hemsley.*Oreodaphne tampicensis*, Meissn. in DC. Prodr. xv. 1, p. 136.SOUTH MEXICO, Tampico (*Berlandier*).10. **Ocotea umbrosa**, Mart., ex Meissn. in DC. Prodr. xv. 1, p. 122.*Oreodaphne umbrosa*, Meissn. loc. cit.

BRAZIL.

 β . ? **bullata**, Meissn. loc. cit.CENTRAL AMERICA (*Ersted*).

5. NECTANDRA.

Nectandra, Roland. in Rottb. Descr. Pl. Surin. p. 10; Benth. et Hook. Gen. Plant. iii. p. 159.

An exclusively American genus of about seventy species, ranging from Mexico and the West Indies to Peru and Brazil.

1. **Nectandra amazonum**, Nees, Syst. Laur. p. 282; DC. Prodr. xv. 1, p. 150 (varietates).PANAMA, Frijoli station (*S. Hayes*, 487), woods near the city of Panama (*Seemann*, 556).—Southward to PERU and BRAZIL. Hb. Kew.2. **Nectandra cuspidata**, Nees, Syst. Laur. p. 330; DC. Prodr. xv. 1, p. 159.PANAMA (*Seemann*, 292; *S. Hayes*, 392).—Southward to BRAZIL and in the WEST INDIES.—Hb. Kew.3. **Nectandra glabrescens**, Benth. Bot. Voy. 'Sulphur,' p. 161; DC. Prodr. xv. 1, p. 165.*Ocotea salicifolia*, Hook. et Arn. Bot. Beech. Voy. p. 309, vix H. B. K., ex Benth.SOUTH MEXICO, San Blas to Acapulco (*Hinds*), San Blas to Tepic (*Sinclair*). Hb. Kew.4. **Nectandra laurel**, Kl.; DC. Prodr. xv. 1, p. 146?PANAMA, Lion Hill (*S. Hayes*, 467). Hb. Kew.Colombia to Peru is the range of *N. laurel* as defined above.5. **Nectandra limbata**, Nees in Linnæa, xxi. p. 509; DC. Prodr. xv. 1, p. 156. MEXICO (*Jurgensen*).—PERU.6. **Nectandra mollis**, Nees, Syst. Laur. p. 287; Linnæa, xxi. p. 504.*a. attenuata*, Meissn. in DC. Prodr. xv. 1, p. 148.*Ocotea mollis*, H. B. K. Nov. Gen. et Sp. p. 164.

COLOMBIA.