

3. **Arceuthobium oxycedri**, Bieb. Fl. Taur. Cauc. Suppl. p. 629; D. Oliv. in Journ. Linn. Soc. vii. p. 104.

Arceuthobium americanum, Nutt., ex Engelm. in Bost. Journ. Nat. Hist. vi. p. 214, in nota; Wats. Bot. Calif. ii. p. 106.

WYOMING to OREGON, and southward to COLORADO and CALIFORNIA.—MEXICO, peak of Orizaba 12,000 feet (*Linden*, 540), summit of San Felipe (*Andrieux*, 349).—SOUTH EUROPE; WEST ASIA. Hb. Kew.

It is uncertain whether *Andrieux* and *Linden*'s specimens be correctly referred to this species, whatever limitation we accept. *Engelmann* regards the American form as a species distinct from *A. oxycedri*.

4. **Arceuthobium**, sp.

NORTH MEXICO, Sierra Madre (*Seemann*, 2138). Hb. Kew.

3. DENDROPHTHORA.

Dendrophthora, Eichl. in Mart. Fl. Bras. v. 2, p. 102, t. 31; Benth. et Hook. Gen. Plant. iii. p. 214.

About twenty-five species are known, most of which inhabit the West Indies, and the remainder continental America.

1. **Dendrophthora biserrula**, Eichl. in Mart. Fl. Bras. v. 2, p. 104.

GUATEMALA (*Friedrichsthal*); COSTA RICA (*Hoffmann*).—COLOMBIA. Hb. Kew.

4. PHORADENDRON.

Phoradendron, Nutt. in Journ. Acad. Philad. ser. 2, i. p. 185; Benth. et Hook. Gen. Plant. iii. p. 214.

An exclusively American genus of about eight species, ranging from Oregon to the Argentine Republic.

1. **Phoradendron angustifolium**, Eichl. in Mart. Fl. Bras. v. 2, p. 115, in obs. ad. n. 13.

Loranthus angustifolius, H. B. K. Nov. Gen. et Sp. iii. p. 442.

Viscum falcatum, Cham. et Schl. in Linnæa, v. p. 172, nec Hook. Ic. t. 368.

Viscum schiedeanaum, DC. Prodr. iv. p. 670.

SOUTH MEXICO, Jalapa (*Linden*, 538; *Schiede & Deppe*), around Oaxaca (*Andrieux*, 346); GUATEMALA (*Friedrichsthal*); COSTA RICA (*Hoffmann*).—PERU. Hb. Kew.

2. **Phoradendron annulatum**, D. Oliv. in Vidensk. Meddel. 1864, p. 176.
COSTA RICA (*Ersted*).

3. **Phoradendron bolleanum**, Hemsley.

Viscum bolleanum, Seem. Bot. Voy. 'Herald,' p. 295, t. 63.

NORTH MEXICO, Sierra Madre (*Seemann*, 2139). Hb. Kew.