

three others may be expected to occur within our limits, as they have been found in the southern part of New Mexico and Arizona.

[i. p. 378.] *Rosa montezumæ*, H. B. K.

S. Watson, loc. cit. p. 351, states that there can be little doubt that this is a variety of *R. canina*, Linn., which exists only as a naturalized plant in North America.

[i. p. 380.] 1. **Cotoneaster denticulata**, H. B. K.; S. Wats. in Proc. Am. Acad. xvii. p. 354.

NORTH MEXICO, in the Sierra Madre, south of Saltillo (*Palmer*, 330).

[i. p. 380.] 1\*. **Cotoneaster nervosa**, Dcne. in Nouv. Archives du Mus. x. p. 177.

MEXICO.

### SAXIFRAGACEÆ.

[i. p. 384.] 4. **Philadelphus serpyllifolius**, A. Gr.; S. Wats. in Proc. Am. Acad. xvii. p. 354.

NORTH MEXICO, in the Sierra Madre, south of Saltillo (*Palmer*).

### CRASSULACEÆ.

[i. p. 392.] 34\*. **Cotyledon schaffneri**, S. Wats. in Proc. Am. Acad. xvii. p. 354.

NORTH MEXICO, mountains around San Luis Potosi (*Schaffner*, 768).

[i. p. 398.] 17\*. **Sedum palmeri**, S. Wats. in Proc. Am. Acad. xvii. p. 355.

NORTH MEXICO, Guajuco, Nuevo Leon (*Palmer*, 2121).

[i. p. 399.] 20\*. **Sedum stelliforme**, S. Wats. in Proc. Am. Acad. xx. p. 364.

NEW MEXICO.—NORTH MEXICO, Huachuca Mountains (*Lemmon*).

### HALORAGÆÆ.

[i. p. 401.] 1. **Callitriche heterophylla**, Pursh, Fl. Am. Sept. i. p. 3; A. Gr. Manual, ed. 5, p. 429; S. Wats. in Proc. Am. Acad. xviii. p. 154.

*Callitriche asagræi*, Hegelm. Monogr. Callit. p. 54, t. 3. fig. 9, t. 4. fig. 1.

NEW YORK and ILLINOIS southward.—NORTH MEXICO, near Morales, San Luis Potosi (*Schaffner*). Hb. Kew.

[i. p. 401.] 1\*. **Callitriche austini**, Engelm.; A. Gr. Man. Bot. Northern U. S. ed. 5, p. 428; S. Wats. in Proc. Am. Acad. xviii. p. 154.

NEW YORK to ILLINOIS and MISSOURI southward.—NORTH MEXICO, near Morales, San Luis Potosi (*Schaffner*).—And in SOUTH AMERICA.