

[ii. p. 164.] 2*. **Sclerocarpus kerberi**, Fournier in Bull. Soc. Bot. France, Comptes Rendus des Séances, xxx. (1883) p. 183.

SOUTH MEXICO, Cordova (*Kerber*).

[ii. p. 166.] 19*. **Montanoa thomasii**, Klatt in Abhandl. naturf. Gesellsch. Halle, xv. (reprint, p. 8).

SOUTH MEXICO, Orizaba (*Thomas*, 1866).

[ii. p. 172.] 2. **Zexmenia brevifolia**, A. Gr.; S. Wats. in Proc. Am. Acad. xviii. p. 105.

TEXAS.—NORTH MEXICO, Soledad, Coahuila (*Palmer*, 625).

[ii. p. 174.] 20. **Zexmenia scandens**, Hemsl.

The specific name was given to this in consequence of Bourgeau designating it a "liane;" but Dr. A. Gray sends a specimen of it to Kew collected by himself at Cordova, with the observation "frutex arborescens valde patens, nec scandens."

[ii. p. 174.] 23. **Zexmenia stenantha**, Hemsl., = *Z. crocea*, A. Gr., fide Gray in Proc. Am. Acad. xix. p. 11.

ii. p. 176, nine lines from bottom, for *recurrens* read *decurrens*.

[ii. p. 177.] 2. **Viguiera canescens**, DC.; S. Wats. in Proc. Am. Acad. xviii. p. 105.

NORTH MEXICO, San Lorenzo de Laguna, Coahuila (*Palmer*, 618).

[ii. p. 178.] 5*. **Viguiera ghiesbreghtii**, A. Gr. in Proc. Am. Acad. xix. p. 6.
SOUTH MEXICO, pine-forests in Morelia (*Ghiesbreght*, 381).

[ii. p. 178.] 7. **Viguiera helianthoides**, H. B. K. vide A. Gr. in Proc. Am. Acad. xix. p. 6.

[ii. p. 178.] 14. **Viguiera seemanni**, Schultz Bip., = *Oyedæa seemanni*, A. Gr. in Proc. Am. Acad. xix. p. 10.

[ii. p. 179.] 1*. **Helianthus annuus**, Linn. Sp. Pl. ed. 2, p. 1276; DC. Prodr. v. p. 585; S. Wats. in Proc. Am. Acad. xviii. p. 105.
Helianthus lenticularis, Dougl. in Bot. Reg. t. 1265.

CALIFORNIA to NEBRASKA and TEXAS.—NORTH MEXICO, without locality (*Palmer*, 600).

[ii. p. 179.] 2. **Helianthus cernuus**, Benth. et Hook., is a true *Flourensia*, vide A. Gr. in Proc. Am. Acad. xix. p. 7.

[ii. p. 179.] 3. **Helianthus ciliaris**, DC., var., S. Wats. in Proc. Am. Acad. xviii. p. 106.

NORTH MEXICO, Juraz, Coahuila (*Palmer*, 616).