

ERICACEÆ.

[ii. p. 283.] 2. **Chimaphila umbellata**, Nutt.

GUATEMALA (*Bernoulli & Cario*, 1964). Hb. Kew.

PLUMBAGINEÆ.

[ii. p. 287.] 1. **Plumbago pulchella**, Boiss.; S. Wats. in Proc. Am. Acad. xviii. p. 112.

NORTH MEXICO, region of San Luis Potosi (*Parry & Palmer*, 240).

Several other specimens enumerated by us under *P. scandens* also belong to this species, thus:—*Botteri*, 810; Guanajuato, *Hartweg*; and *Galeotti*, 1757, in part.

[ii. p. 287.]

PRIMULACEÆ.

The now widely diffused *Anagallis arvensis*, Linn., was collected at Saltillo, by Palmer, and San Luis Potosi, by Schaffner; S. Wats. in Proc. Am. Acad. xviii. p. 112.

Gray now (*Coulter's Bot. Gaz.* 1886, p. 231), in a revision of *Dodecatheon*, defines five species.

ii. p. 299, fifteen lines from top, for *subsessiflora* read *subsessiliflora*.

SAPOTACEÆ.

[ii. p. 299.] 9. **Bumelia spinosa**, DC.; S. Wats. in Proc. Am. Acad. xviii. p. 112.

ARIZONA; TEXAS.—NORTH MEXICO, Saltillo, Coahuila (*Palmer*, 786).

EBENACEÆ.

[ii. p. 300.] 4. **Diospyros texana**, Scheele; S. Wats. in Proc. Am. Acad. xviii. p. 112.

NORTH MEXICO, Guajuco, Nuevo Leon (*Palmer*, 784).

OLEACEÆ.

[ii. p. 304.] 2. **Menodora helianthemoides**, Humb. et Bonpl.; S. Wats. in Proc. Am. Acad. xviii. p. 112.

Watson regards Parry & Palmer's 570 as a small-flowered form of this species rather than *M. coulteri*, A. Gr.

[ii. p. 304.] 4. **Menodora longiflora**, A. Gr.; S. Wats. in Proc. Am. Acad. xviii. p. 112.

NORTH MEXICO, Monclova and Juraz, Coahuila (*Palmer*, 792, 794).

[ii. p. 304.] 6. **Menodora scabra**, A. Gr.; S. Wats. in Proc. Am. Acad. xviii. p. 304.

NORTH MEXICO, at Juraz on the Sabinas river, Coahuila (*Palmer*, 793).