

APOCYNACEÆ.

[ii. p. 307.] 1. **Rauwolfia heterophylla**, Rœm. et Schult.
GUATEMALA (*Bernoulli & Cario*, 1831). Hb. Kew.

[ii. p. 307.] 2. **Rauwolfia longifolia**, A. DC.
GUATEMALA (*Bernoulli & Cario*, 1809). Hb. Kew.

[ii. p. 313.] 2. **Apocynum cannabinum**, Linn.; S. Wats. in Proc. Am. Acad. xviii. p. 113.

Palmer's 806, referred by us to *A. androsæmifolium*, is this species, and the latter is not known to occur in Mexico.

[ii. p. 314.] 3*. **Echites (Euechites) coulteri**, S. Wats. in Proc. Am. Acad. xviii. p. 113.

NORTH MEXICO, south of Saltillo (*Palmer*, 805), without locality (*Coulter*, 987).

ASCLEPIADEÆ.

[ii. p. 322.] 2. *Acerates circinalis*, *A. asperula*, and *A. gomphocarpoides* are referred by Fournier (Ann. Sc. Nat. série 6, xiv. p. 369) to *Asclepiodora*. *Gomphocarpus arachnoideus*, Fourn. in Bull. Soc. Bot. France, xiv. p. 250, raised at Paris from Mexican seeds, we had overlooked, as well as *G. hypoleucus*, A. Gray, from the Santa Rita Mountains.

[ii. p. 322.] 4*. **Acerates viridiflora**, Ell.; Chapm. Fl. Southern U.S. p. 365; S. Wats. in Proc. Am. Acad. xviii. p. 114.

EASTERN NORTH AMERICA.—NORTH MEXICO, Caracol Mountains, Coahuila (*Palmer*).

[ii. p. 322.] 4**. **Acerates schaffneri**, Fournier in Ann. Sc. Nat. série 6, xiv. p. 386.

MEXICO, without locality (*Schaffner*, 861).

[ii. p. 322.] 4***. **Acerates vinosa**, Fournier in Ann. Sc. Nat. série 6, xiv. p. 387.

SOUTH MEXICO, among pines at an elevation of 9000 feet (*Ghiesbreght*, 35).

[ii. p. 322.] **Asclepias alticola**, Fournier in Ann. Sc. Nat. série 6, xiv. p. 371.

SOUTH MEXICO, in the cordillera between Cuernavaca and Toluca (*Ghiesbreght*).

[ii. p. 323.] 7. *Asclepias elata*, Benth., = **A. glaucescens**, H. B. K., fide S. Wats. in Proc. Am. Acad. xviii. p. 114.

[*Asclepias galeottii*, Fourn. in Bull. Soc. Bot. France, xiv. p. 250 (*Blepharodon*, Galeotti), not taken up by the author in his review of the genus in the *Annales*.]