

[ii. p. 432.] 24. **Cestrum multinervium**, Dun. ; S. Wats. in Proc. Am. Acad. xviii. p. 128.

TEXAS.

[ii. p. 433.] 26. *Cestrum foetidissimum*, Jacq. Hort. Schœnb. iii. t. 329, is specifically distinct from **C. nocturnum**, to which we, following Grisebach (Fl. Brit. W. Ind. p. 444), had referred it; and it is probably not found within our area.

[ii. p. 434.] 40\*. **Cestrum warscewiczii**, Klotzsch in Otto & Dietr. Allg. Gart. Zeit. xix. p. 362.

COSTA RICA, Cartago (*Warscewicz*).

Closely allied to *C. aurantiacum*, Lindl., and originally known in the gardens of Europe under the name of *Habrothamnus aureus*.

[ii. p. 435.] 4\*. **Nicotiana nudicaulis**, S. Wats. in Proc. Am. Acad. xviii. p. 128.

NORTH MEXICO, Monterey, Nuevo Leon (*Palmer*, 961).

[ii. p. 435.] 12\*. **Nicotiana trigonophylla**, Dun. in DC. Prodr. xi. p. 562 ; A. Gr. Synopt. Fl. N. Am. ii. p. 242 ; S. Wats. in Proc. Am. Acad. xviii. p. 128.

CALIFORNIA to TEXAS.—NORTH MEXICO, in the mountains east of Saltillo, Coahuila (*Palmer*, 962), in rocky places about San Luis Potosi (*Schaffner*, 688 ; *Parry & Palmer*, 660).

ii. p. 438, top line, for *Brachylossis* read *Brachyglossis*.

## SCROPHULARINEÆ.

[ii. p. 444.] 5. **Pentstemon campanulatus**, Willd. ; S. Wats. in Proc. Am. Acad. xviii. p. 129.

NORTH MEXICO, at Lerios, in the high mountains east of Saltillo, Coahuila (*Palmer*, 976).

[ii. p. 445.] 15\*. **Pentstemon rubescens**, A. Gr. in Proc. Am. Acad. xix. p. 92. NORTH MEXICO, in Tanner's Cañon, near Fort Huachuca (*Lemmon*).

[ii. p. 449.] 2. **Mimulus glabratus**, H. B. K. ; S. Wats. in Proc. Am. Acad. xviii. p. 130.

NORTH MEXICO, at Saltillo and San Lorenzo de Laguna (*Palmer*, 977, 979).

[ii. p. 451.] 1. **Herpestis chamædryoides**, H. B. K. ; S. Wats. in Proc. Am. Acad. xviii. p. 130.

Watson states that the form we have referred to *H. nigrescens*, Benth., is really this species.