

lobis labii inferioris brevioribus obtusioribus. Staminibus longius exsertis; stylo longitudinem staminum subæquante parum curvato."

The specimens in Kew Herbarium from Jalapa (*Linden*, 198) and Vera Cruz to Orizaba (*Müller*, 647) answer well to the foregoing description as to slenderness, and in being cylindrical to the base.

LENTIBULARIACEÆ.

[ii. p. 469.] 4. **Utricularia endresii**, Reichb. f. Bot. Mag. t. 6656.

[ii. p. 471.] 2*. **Pinguicula sodalium**, Fourn. in Bull. Soc. Bot. France, xx. p. lxvii.

SOUTH MEXICO, Orizaba (*Müller*, 114).

This probably is the same as *P. caudata*.

GESNERACEÆ.

[ii. p. 473.] 3*. **Achimenes cardinalis**, A. Dietr. in Otto & Dietr. Allg. Gart. Zeit. xv. p. 314.

MEXICO.

[ii. p. 480.] 1*. **Gesnera rugata**, Scheidw. in Otto & Dietr. Allg. Gart. Zeit. xv. p. 226.

MEXICO (*Ghiesbreght*).

ii. p. 480, eight lines from top, for *tubiflora* read *tubiflorum*.

BIGNONIACEÆ.

[ii. p. 491.] 15*. **Bignonia peltata**, Otto & Dietr. Allg. Gart. Zeit. ix. p. 58.

MEXICO.

ACANTHACEÆ.

[ii. p. 500.] *Elytraria bromoides*, CErst.; S. Wats. in Proc. Am. Acad. xviii. p. 133.

Watson identifies Palmer's 2029 (Guajuco, Nuevo Leon) with this, which he says is quite distinct from *E. tridentata*, Vahl, to which we had referred it.

[ii. p. 503.] 6. **Calophanes jasminum-mexicanum**, Nees; S. Wats. in Proc. Am. Acad. xviii. p. 133.

Watson suggests that *C. decumbens*, A. Gr., and *C. schiedeana*, Nees, are the same as this.