

		feet.
<i>Cheilanthes lendigera</i> (N. & S.)	Vaqueria del Jacal, Mexico.	10,000
— <i>microphylla</i> (N. & S.)	Oaxaca, Mexico.	8,000
— <i>scariosa</i> (A.)	Orizaba, „	12,000
— <i>speciosissima</i> (E.)	Orizaba, „	10,000 to 12,000
<i>Asplenium</i> (W.) <i>filix-fœmina</i> (W.)	Orizaba, „	9,500 to 12,000
— <i>monanthemum</i> (W.)	Mexico.	1,000 to 11,500
— <i>trichomanes</i> (W.)	Mexico.	to 11,000
	Alotenango, Mexico.	above 11,000
	Fuego, Guatemala.	above 11,000
<i>Nephrodium</i> (W.) <i>sphærocarpum</i> (E.)	Real del Monte, Mexico.	8,000
<i>Polypodium</i> (W.) <i>heteromorphum</i> (A.)	Orizaba, „	10,000 to 12,000
— <i>lanceolatum</i> (W.)	Orizaba, „	9,000 to 12,000
— <i>moniliforme</i> (A.)	Orizaba, „	9,000 to 12,000
— <i>plebeium</i> (S.)	Coban, Guatemala.	8,000 to 10,000
	Fuego, „	8,000 to 10,000
— <i>rigescens</i> (W.)	Cerro de Pelado, Mexico.	7,000 to 10,000
<i>Notholaena</i> (W.) <i>aschenborniana</i> (N.)	Oaxaca, „	8,000
<i>Gymnogramme</i> (W.) <i>pilosa</i> (E.)	Orizaba, „	9,760 to 10,000
<i>Antrophyum</i> (W.) <i>ensiforme</i> (E.)	Totontepec, „	10,000
<i>Acrostichum</i> (W.) <i>lepidotum</i> (S.)	Orizaba, „	10,000 to 11,000
— <i>lindenii</i> (S.)	Orizaba, „	9,000 to 10,000
— <i>mathewsii</i> (A.)	Orizaba, „	9,500 to 11,000
— <i>simplex</i> (S.)	Orizaba, „	10,000
— <i>squamipes</i> (S.)	Sempoaltepec, „	8,000 to 9,000
— <i>tectum</i> (S.)	Chinantla, „	7,000 to 8,000
<i>Marattia</i> (W.) <i>alata</i> (S.)	Oaxaca, „	7,000 to 8,000
<i>Lycopodium</i> (W.) <i>clavatum</i> (W.)	Sempoaltepec, „	11,000 to 12,000

Analysis of the Distribution of the Genera and Species in the foregoing list.

GENERA.					
Endemic.	Extending to				
	North America.	South America.	Andes only.	N. and S. America.	Beyond America.
13	9	25	22	26	165

America ..... 82 = 31.6 per cent.

Endemic ..... 13 = 5.0 „

Of wider range .. 165 = 63.4 „

Total ..... 260 100.0