

HELICINA LACINIOSA (*Mighels*).

*Testa parva, conico-globosa, solidula, cinnamomea lineis flexuosis luteis variegata: spira anfractibus quinque subplanulatis, ultimo angulato et striis elevatis duobus ad tribus cincto: apertura lunata; labro acuto, ad columellam expanso; operculo tenui, fulvo.*

*Helicina laciniosa*, MIGHELS; Proceed. Boston Soc. Nat. Hist., ii. 19. Jan. 1845.

SHELL conic-globose, rather solid, of a cinnamon-brown colour, variegated with yellowish zigzag stripes. Spire composed of five flattened whorls, the last of which is obtusely angular at periphery, and has two or three raised revolving lines upon it. The aperture is lunate, the lip acute, expanding in the vicinity of the columella. Operculum thin, yellowish-brown.

Axis one-tenth of an inch; base three-twentieths of an inch.

Found at Kauai, Sandwich Islands. *Couthouy*. Waianai, *Pickering*.

Resembles the two preceding; but is distinguished from *H. fulgora* by being smaller and more globular, and from *H. musiva*, by the raised revolving lines, and by having zigzag markings both on the base and on the spire.

Figures 108, 108 *a*, 108 *b*, three views of the shell, magnified three diameters; 108 *c*, natural size.

HELICINA TROCHLEA (*Gould*).

*Testa minuta, trochiformis, supra cinerea, costis acutis volventibus tribus vel quatuor (intervallis concavis, scabrosis) cincta, infra convexa, polita, citrina: spira anfractibus quatuor, ultimo ad peripheriam bicarinato: apertura sub-trigona; peristomate leviter reflexo; columellâ ad basim angulatâ, callo copioso indutâ.*